

UNITED STATES PATENT APPLICATION

of

Joseph P. Steiner
and
Gregory S. Hamilton

for

ROTAMASE ENZYME ACTIVITY INHIBITORS

Finnegan, Henderson, Farabow, Garrett & Dunner
1300 I Street, N.W., Suite 700
Washington, D.C. 20005
Telephone: (202) 408-4000
Facsimile: (202) 408-4400

This application is a continuation-in-part of U.S. Application Serial No. 08/551,026, filed October 31, 1995, and of U.S. Application Serial No 09/359,351, filed July 21, 1999, which is a continuation of U.S. Application Serial No. 08/693,003, filed August 6, 1996, which is a continuation of U.S. Application Serial No 08/479,436, filed June 7, 1995, now U.S. Patent No. 5,614,547, which are hereby incorporated by reference in their entirety.

This invention relates to neurotrophic compounds having an affinity for FKBP-type immunophilins, their preparation and use as inhibitors of the enzyme activity associated with immunophilin proteins, and particularly inhibitors of peptidyl-prolyl isomerase or rotamase enzyme activity, and their use as small molecule neurotrophic drugs.

The term immunophilin refers to a number of proteins that serve as receptors for the principal immunosuppressant drugs, cyclosporin A (CsA), FK506, and rapamycin. Known classes of immunophilins are cyclophilins and FK506 binding proteins, such as FKBP. Cyclosporin A binds to cyclophilin while FK506 and rapamycin bind to FKBP. These immunophilin-drug complexes interface with a variety of intracellular signal transduction systems, especially in the immune system and the nervous system.

Immunophilins are known to have peptidyl-prolyl isomerase (PPIase) or rotamase enzyme activity. It has been determined that rotamase activity has a role in the catalyzation of the interconversion of the cis and trans isomer of the substrate proteins of the immunophilin.

Immunophilins were originally discovered and studied in immune tissue. It was initially postulated by those skilled in the art that inhibition of the immunophilin's

rotamase activity leads to the inhibition of T-cell proliferation, thereby causing the immunosuppressive action exhibited by immunosuppressive drugs such as cyclosporin A, FK506, and rapamycin. Further study has shown that the inhibition of rotamase activity, in and of itself, is not sufficient for immunosuppressant activity. Instead

5 immunosuppression appears to stem from the formation of a complex of immunosuppressant drugs and immunophilins. It has been shown that the immunophilin-drug complexes interact with ternary protein targets as their mode of action. In the case of FKBP-FK506 and cyclophilin-CsA, the drug-immunophilin complexes bind to the enzyme calcineurin, inhibiting T-cell receptor signaling leading to

10 T-cell proliferation. Similarly, the complex of rapamycin and FKBP interacts with the RAFT1/FRAP protein and inhibits signaling from the IL-2 receptor.

Immunophilins have been found to be present at high concentrations in the central nervous system. Immunophilins are enriched 10-50 times more in the central nervous system than in the immune system. Within neural tissues, immunophilins

15 appear to influence nitric oxide synthesis, neurotransmitter release, and neuronal process extension.

FK506 also augments the phosphorylation of growth-associated protein-43 (GAP43). GAP43 is involved in neuronal process extension and its phosphorylation appears to augment this activity. Accordingly, the effects of FK506, rapamycin, and

20 cyclosporin in neuronal process extension have been examined using PC12 cells. PC12 cells are a continuous line of neuronal-like cells which extend neurites when stimulated by nerve growth factor (NGF).

Surprisingly, it has been found that picomolar concentrations of an

immunosuppressant such as FK506 or rapamycin stimulate neurite outgrowth in PC12 cells and sensory neurons, namely dorsal root ganglion cells (DRGs). In whole animal experiments, FK506 has been shown to stimulate nerve regeneration following facial nerve injury and results in functional recovery in animals with sciatic nerve lesions.

5 More particularly, it has been found that drugs with a high affinity for FKBP are potent rotamase inhibitors and exhibit excellent neurotrophic effects. Snyder et al., "Immunophilins and the Nervous System", *Nature Medicine*, Volume 1, No. 1, January 1995, 32-37. These findings suggest the use of immunosuppressants in treating various peripheral neuropathies and in enhancing neuronal regrowth in the central
10 nervous system (CNS). Studies have demonstrated that neurodegenerative disorders such as senile dementia of the Alzheimer's type (Alzheimer's disease, SDAT), Parkinson's disease, and amyotrophic lateral sclerosis (ALS) may occur due to the loss, or decreased availability, of a neurotrophic substance specific for a particular population of neurons affected in the disorder.

15 Several neurotrophic factors effecting specific neuronal populations in the central nervous system have been identified. For example, it has been hypothesized that Alzheimer's disease results from a decrease or loss of nerve growth factor (NGF). It has thus been proposed to treat SDAT patients with exogenous NGF or other neurotrophic proteins such as brain derived growth factor (BDNF), glial derived growth
20 factor, ciliary neurotrophic factor (CNTF), and neurotrophin-3 (NT-3) to increase the survival of degenerating neuronal populations.

Clinical application of these proteins in various neurological disease states is hampered by difficulties in the delivery and bioavailability of large proteins to nervous

system targets. By contrast, immunosuppressant drugs with neurotrophic activity are relatively small and display specificity. However, when administered chronically, immunosuppressants exhibit a number of potentially serious side effects including nephrotoxicity, such as impairment of glomerular filtration and irreversible interstitial fibrosis (Kopp et al., 1991, *J. Am. Soc. Nephrol.* 1:162); neurological deficits, such as involuntary tremors, or non-specific cerebral angina such as non-localized headaches (De Groen et al., 1987, *N. Engl. J. Med.* 317:861); and vascular hypertension with complications resulting therefrom (Kahan et al., 1989 *N. Engl. J. Med.* 321: 1725).

The present invention provides non-immunosuppressive neurotrophic compounds having an affinity for FKBP-type immunophilins that are extremely potent in augmenting neurite outgrowth, for promoting neuronal growth, and for facilitating regeneration in various neuropathological situations where neuronal repair can be facilitated. Such neuropathological situations include peripheral nerve damage by physical injury or disease state such as diabetes, physical damage to the central nervous system (spinal cord and brain), brain damage associated with stroke, and neurological disorders relating to neurodegeneration, including Parkinson's disease, Alzheimer's disease, and amyotrophic lateral sclerosis.

SUMMARY OF THE INVENTION

This invention relates to neurotrophic compounds having an affinity for FKBP-type immunophilins and to methods of using neurotrophic compounds having an affinity for FKBP-type immunophilins.

One embodiment of this invention is neurotrophic compounds of the formula I, detailed below.

Another embodiment of this invention is neurotrophic compounds of the formula II, detailed below.

Another embodiment of this invention is a method of treating a neurological activity in an animal, comprising: administering to an animal an effective amount of a
5 neurotrophic compound having an affinity for FKBP-type immunophilins to stimulate growth of damaged peripheral nerves or to promote neuronal regeneration, wherein the FKBP-type immunophilin exhibits rotamase activity.

Another embodiment of this invention is a method of treating a neurological disorder in an animal, comprising: administering to an animal an effective amount of a
10 neurotrophic compound having an affinity for FKBP-type immunophilins in combination with an effective amount of a neurotrophic factor selected from the group consisting of neurotrophic growth factor, brain derived growth factor, glial derived growth factor, ciliary neurotrophic factor, and neurotrophin-3, to stimulate growth of damaged peripheral nerves or to promote neuronal regeneration, wherein the FKBP-type immunophilin
15 exhibits rotamase activity.

Another embodiment of this invention is a method of stimulating growth of damaged peripheral nerves, comprising: administering to damaged peripheral nerves an effective amount of a neurotrophic compound having an affinity for FKBP-type immunophilins to stimulate or promote growth of the damaged peripheral nerves,
20 wherein the FKBP-type immunophilins exhibit rotamase activity.

Another embodiment of this invention is a method for promoting neuronal regeneration and growth in animals, comprising: administering to an animal an effective amount of a neurotrophic compound having an affinity for FKBP-type immunophilins to

promote neuronal regeneration, wherein the FKBP-type immunophilins exhibit rotamase activity.

Yet another embodiment of this invention is a method for preventing neurodegeneration in an animal, comprising: administering to an animal an effective
 5 amount of a neurotrophic compound having an affinity for FKBP-type immunophilins to prevent neurodegeneration, wherein the FKBP-type immunophilins exhibit rotamase activity.

DETAILED DESCRIPTION OF THE INVENTION


The novel neurotrophic compounds of this invention are relatively small
 10 molecules in relation to other known compounds, such as rapamycin, FK506, and cyclosporin.

The neurotrophic compounds of this invention have an affinity for the FK506 binding proteins such as FKBP-12. When the neurotrophic compounds of the invention are bound to FKBP, they have been found to inhibit the prolyl-peptidyl cis-trans
 15 isomerase activity, or rotamase activity of the binding protein. The compounds of the invention also have been found to stimulate neurite growth, while not exhibiting an immunosuppressive effect. That is, the compounds of the invention are non-immunosuppressive.

The term "non-immunosuppressive" refers to the inability of the compounds of
 20 the present invention to suppress the immune system when compared to a control such as FK506 or cyclosporin A. Assays for determining whether a compound is immunosuppressive are well known to those of ordinary skill in the art. Specific non-limiting examples of well known assays include PMA and OKT3 assays wherein

mitogens are used to stimulate proliferation of human peripheral blood lymphocytes (PBC). Compounds added to such assay systems are evaluated for their ability to inhibit such proliferation.

In one embodiment, this invention relates to a novel class of neurotrophic
5 compounds represented by the formula I:




Formula I

and pharmaceutically acceptable salts thereof,

- wherein Y is CH₂, O, NH, or N-(C1-C4 alkyl);

10 - wherein Z and R₂ are independently Ar, (C5-C7)-cycloalkyl substituted (C1-C6)-straight or branched alkyl or alkenyl, (C5-C7)-cycloalkenyl substituted (C1-C6)-straight or branched alkyl or alkenyl, or Ar substituted (C1-C6)-straight or branched alkyl or alkenyl, wherein in each case, one or two carbon atoms of the straight or branched alkyl or alkenyl groups may be substituted with 1-2

15 heteroatoms selected from the group consisting of oxygen, sulfur, SO and SO₂ in chemically reasonable substitution patterns, or



- wherein Q is hydrogen, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl;

5 - wherein T is Ar or substituted 5-7 membered cycloalkyl with substituents at positions 3 and 4 which are independently selected from the group consisting of hydrogen, hydroxyl, O-(C1-C4)-alkyl or O-(C1-C4)-alkenyl and carbonyl;

 - wherein Ar is selected from the group consisting of monocyclic and bicyclic heterocyclic aromatic ring systems with individual ring sizes being 5 or 6 which may contain in either or both rings a total of 1-4 heteroatoms
10 independently selected from oxygen, nitrogen and sulfur; wherein 1-naphthyl, 2-naphthyl, 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl and phenyl are preferred, and wherein Ar may contain one to three substituents which are independently selected from the group consisting of hydrogen, halo,
15 hydroxyl, hydroxymethyl, nitro, CF₃, trifluoromethoxy, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, O-benzyl, O-phenyl, amino, 1,2-methylenedioxy, carbonyl and phenyl;

 - wherein R₁ is either hydrogen or U; X is either oxygen or CH-U, provided that if R₁ is hydrogen, then X is CH-U, or if X is oxygen then R₁ is U;

20 - wherein U is hydrogen, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl, (C5-C7)-cycloalkyl, (C5-C7)-cycloalkenyl substituted with (C1-C4)-straight or branched alkyl or (C1-C4)-straight or branched alkenyl, [(C1-C4)-alkyl or (C1-C4)-alkenyl]-Ar or Ar (Ar as described above);


- wherein J is hydrogen or C1 or C2 alkyl or benzyl; K is (C1-C4)-straight or branched alkyl, benzyl or cyclohexylethyl; or wherein J and K may be taken together to form a 5-7 membered heterocyclic ring which may contain an oxygen (O), sulfur (S), SO or SO₂ substituted therein; and

5 wherein n is 0-3.

The stereochemistry at position 1 (Formula I) is (R) or (S), with (S) preferred.

The stereochemistry at position 2 is (R) or (S).

In a second embodiment, a novel class of neurotrophic compounds of this invention are represented by the formula II:




Formula II

10

and pharmaceutically acceptable salts thereof,

- wherein Y is O, NH, or N-(C1-C4 alkyl);

- wherein Z is hydrogen, CHL-Ar, (C1-C6)-straight or branched alkyl, (C1-C6)-straight or branched alkenyl, (C5-C7)-cycloalkyl, (C5-C7)-cycloalkenyl or Ar
15 substituted (C1-C6)-alkyl or alkenyl, or



- wherein L and Q are independently hydrogen, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl;

5 - wherein T is Ar or substituted cyclohexyl with substituents at positions 3 and 4 which are independently selected from the group consisting of hydrogen, hydroxyl, O-(C1-C4)-alkyl or O-(C1-C4)-alkenyl and carbonyl;

10 - wherein Ar is selected from the group consisting of 1-naphthyl, 2-naphthyl, 2-furyl, 3-furyl, 2-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl and phenyl having one to three substituents which are independently selected from the group consisting of hydrogen, halo, hydroxyl, nitro, CF₃, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, O-benzyl, O-phenyl, amino and phenyl;

15 - wherein R₁ is either hydrogen or U; X is either oxygen or CH-U, provided that if R₁ is hydrogen, then X is CH-U, or if X is oxygen then R₁ is U;

20 - wherein U is hydrogen, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, C1-C6-straight or branched alkyl, Ar or C1-C6-straight or branched alkenyl, C5-C7-cycloalkyl, (C5-C7)-cycloalkenyl substituted with (C1-C4)-straight or branched alkyl or (C1-C4)-straight or branched alkenyl, 2-indolyl, 3-indolyl, [(C1-C4)-alkyl or (C1-C4)-alkenyl]-Ar or Ar (Ar as described above);

- wherein J is hydrogen or C1 or C2 alkyl or benzyl; K is (C1-C4)-straight

or branched alkyl, benzyl or cyclohexylethyl; or wherein J and K may be taken together to form a 5-7 membered heterocyclic ring which may contain an oxygen (O), sulfur (S), SO or SO₂ substituted therein.

The stereochemistry at position 1 (Formula II) is (R) or (S), with (S) preferred.

5 The compounds of this invention exist as stereoisomeric forms, either as enantiomers or diastereoisomers. The stereochemistry at position 1 of Formula I or II is R or S, with S preferred. Included within the scope of the invention are the enantiomers, the racemic form, and the diastereoisomeric mixtures. Enantiomers as well as diastereoisomers can be separated by methods known to those skilled in the art.

10 It is known that immunophilins such as FKBP preferentially recognize peptide substrates containing Xaa-Pro-Yaa motifs, where Xaa and Yaa are lipophilic amino acid residues. Schreiber et al. 1990 *J. Org. Chem.* 55, 4984-4986; Harrison and Stein, 1990 *Biochemistry*, 29, 3813-3816. Thus, modified prolyl peptidomimetic compounds bearing lipophilic substituents should bind with high affinity to the hydrophobic core of the FKBP
15 active site and inhibit its rotamase activity.

 The compounds of the present invention can be used in the form of salts derived from inorganic or organic acids and bases. Included among such acid salts are the following: acetate, adipate, alginate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate,
20 dodecylsulfate, ethanesulfonate, fumarate, glucoheptanoate, glycerophosphate, hemissulfate heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, oxalate, pamoate, pectinate, propionate, succinate, tartrate, thiocyanate,

tosylate and undecanoate. Base salts include ammonium salts, alkali metal salts such as sodium and potassium salts, alkaline earth metal salts such as calcium and magnesium salts, salts with organic bases such as dicyclohexylamine salts, N-methyl-D-glucamine, and salts with amino acids such as arginine, lysine, and so forth. Also,

5 the basic nitrogen-containing groups can be quarternized with such agents as lower alkyl halides, such as methyl, ethyl, propyl, and butyl chlorides, bromides and iodides; dialkyl sulfates such as dimethyl, diethyl, dibutyl and diamyl sulfates; long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides; aralkyl halides like benzyl and phenethyl bromides; and others. Water or oil-soluble or

10 dispersible products are thereby obtained.

The neurotrophic compounds of this invention can be periodically administered to a patient undergoing treatment for neurological disorders or for other reasons in which it is desirable to stimulate neuronal regeneration and growth, such as in various peripheral neuropathic and neurological disorders relating to neurodegeneration. The

15 compounds of this invention can also be administered to animals, including mammals other than humans, for treatment of various neurological disorders.

The novel compounds of the present invention are potent inhibitors of rotamase activity and possess an excellent degree of neurotrophic activity. The neurotrophic activity is useful in the stimulation of growth of damaged neurons, the promotion of

20 neuronal regeneration, the prevention of neurodegeneration, and in the treatment of several neurological disorders known to be associated with neuronal degeneration and peripheral neuropathies. The neurological disorders that may be treated include, but are not limited to: trigeminal neuralgia, glossopharyngeal neuralgia, Bell's Palsy,

myasthenia gravis, muscular dystrophy, amyotrophic lateral sclerosis, progressive muscular atrophy, progressive bulbar inherited muscular atrophy, herniated, ruptured or prolapsed intervertebral disk syndromes, cervical spondylosis, plexus disorders, thoracic outlet destruction syndromes, peripheral neuropathic such as those caused by lead, 5 dapsone, ticks, porphyria, or Guillain-Barré syndrome, Alzheimer's disease, and Parkinson's disease.

For these purposes the compounds of the present invention may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir in dosage formulations containing conventional non-toxic 10 pharmaceutically-acceptable carriers, adjuvants and vehicles. The term parenteral as used herein includes subcutaneous, intravenous, intramuscular, intraperitoneal, intrathecal, intraventricular, intrasternal and intracranial injection or infusion techniques.

To be effective therapeutically, the compounds of the invention should readily penetrate the blood-brain barrier when peripherally administered. Compounds of this 15 invention which cannot penetrate the blood-brain barrier can be effectively administered by an intraventricular route.

The pharmaceutical compositions may be in the form of a sterile injectable preparation, for example as a sterile injectable aqueous or oleaginous suspension. This suspension may be formulated according to techniques known in the art using suitable 20 dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution

and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. Fatty acids such as oleic acid and its glyceride derivatives find use in the preparation of injectables, as do olive oil or
5 castor oil, especially in their polyoxyethylated versions. These oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant.

The compounds may be administered orally in the form of capsules or tablets, for example, or as an aqueous suspension or solution. In the case of tablets for oral use, carriers which are commonly used include lactose and corn starch. Lubricating agents,
10 such as magnesium stearate, are also typically added. For oral administration in a capsule form, useful diluents include lactose and dried corn starch. When aqueous suspensions are required for oral use, the active ingredient is combined with emulsifying and suspending agents. If desired, certain sweetening and/or flavoring and/or coloring agents may be added.

15 The compounds of this invention may also be administered in the form of suppositories for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at room temperature but liquid at rectal temperature and therefore will melt in the rectum to release the drug. Such materials include cocoa butter, beeswax and polyethylene
20 glycols.

The compounds of this invention may also be administered topically, especially when the conditions addressed for treatment involve areas or organs readily accessible by topical application, including neurological disorders of the eye, the skin, or the lower

intestinal tract. Suitable topical formulations are readily prepared for each of these areas.

For ophthalmic use, the compounds can be formulated as micronized suspensions in isotonic, pH adjusted sterile saline, or, preferably, as solutions is
 5 isotonic, pH adjusted sterile saline, either with or without a preservative such as benzylalkonium chloride. Alternatively for the ophthalmic uses the compounds may be formulated in an ointment such as petrolatum.

For application topically to the skin, the compounds can be formulated in a suitable ointment containing the compound suspended or dissolved in, for example, a
 10 mixture with one or more of the following: mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene polyoxypropylene compound, emulsifying wax and water. Alternatively, the compounds can be formulated in a suitable lotion or cream containing the active compound suspended or dissolved in, for example, a mixture of one or more of the following: mineral oil, sorbitan monostearate, polysorbate
 15 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

Topical application for the lower intestinal tract can be effected in a rectal suppository formulation (see above) or in a suitable enema formulation.

Dosage levels on the order of about 0.1 mg to about 10,000 mg of the active ingredient compound are useful in the treatment of the above conditions, with preferred
 20 levels of about 0.1 mg to about 1,000 mg. The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration.

It is understood, however, that a specific dose level for any particular patient will

depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, rate of excretion, drug combination, and the severity of the particular disease being treated and form of administration.

- 5 The compounds can be administered with other neurotrophic agents such as neurotrophic growth factor (NGF), glial derived growth factor, brain derived growth factor, ciliary neurotrophic factor, and neurotrophin-3. The dosage level of other neurotrophic drugs will depend upon the factors previously stated and the neurotrophic effectiveness of the drug combination.

10

Methods and Procedures

K_i Test Procedure

Inhibition of the peptidyl-prolyl isomerase (rotamase) activity of the inventive compounds can be evaluated by known methods described in the literature (Harding, M.W. et al. *Nature* 341: 758-760 (1989); Holt et al. *J. Am. Chem. Soc.* 115: 9923-9938).

- 15 These values are obtained as apparent K_i values and are presented in Table I. The *cis-trans* isomerization of an phenylalanine-proline bond in a model substrate, N-succinyl-Ala - Phe -Pro-Phe-*p*-nitroanilide, is monitored spectrophotometrically in a chymotrypsin-coupled assay, which releases *para*-nitroanilide from the *trans* form of the substrate. The inhibition of this reaction caused by the addition of different
- 20 concentrations of inhibitor is determined, and the data is analyzed as a change in first-order rate constant as a function of inhibitor concentration to yield the apparent K_i values.


In a plastic cuvette are added 950 µL of ice cold assay buffer (25 mM HEPES,

1 mM dithiothreitol), 25 μ L of chymotrypsin (50 mg/ml in 1 mM HCl) and 10 μ L of test compound at various concentrations in dimethyl sulfoxide. The reaction is initiated by the addition of 5 μ L of substrate (succinyl-Ala-Phe-Pro-Phe-para-nitroanilide, 5 mg/mL in 2.35 mM LiCl in trifluoroethanol).

- 5 The absorbance at 390 nm versus time is monitored for 90 sec using a spectrophotometer and the rate constants are determined from the absorbance versus time data files.

The data for these experiments are presented in Table I and in Table II.


TABLE I



10

No.	Z	R ₁	m	K _i μ M
15	1 Benzyl	Phenyl	2	1.5
	2 3-Phenylpropyl	Phenyl	2	---
	3 4-(4-Methoxy-phenyl)butyl	Phenyl	2	---
	4 4-Phenylbutyl	Phenyl	2	0.35
	5 Phenethyl	Phenyl	2	1.1
20	6 4-Cyclohexyl-butyl	Phenyl	2	0.4
	7 Benzyl	Methoxy	2	80
	8 4-Cyclohexyl-butyl	Methoxy	2	6
25	9 3-Cyclohexyl-propyl	Methoxy	2	20
	10 3-Cyclopentyl-propyl	Methoxy	2	35
	11 Benzyl	2-Furyl	2	3
30	12 4-Cyclohexyl-	3,4,5-Trimethoxy-	2	0.04

		butyl	phenyl		
	13	3-Phenoxy-benzyl	3,4,5-Trimethoxy-phenyl	2	0.018
5	14	4-Phenylbutyl	3,4,5-Trimethoxy-phenyl	2	0.019
	15	3-(3-Indolyl)propyl	3,4,5-Trimethoxy-phenyl	2	0.017
	16	4-(4-Methoxy-phenyl)butyl	3,4,5-Trimethoxy-phenyl	2	0.013
10	17	3-phenyl-1-propyl	1,1-dimethylpropyl	1	0.042
	18	3-phenyl-1-prop-2-(E)-enyl	1,1-dimethylpropyl	1	0.125
	19	3-(3,4,5-trimethoxy-phenyl)-1-propyl	1,1-dimethylpropyl	1	0.025
15	20	3-(3,4,5-trimethoxy-phenyl)-1-prop-2-(E)-enyl	1,1-dimethylpropyl	1	0.125
	21	3-(4,5-dichloro-phenyl)-1-prop-2-(E)-enyl	1,1-dimethylpropyl	1	2.50
20	22	3-(2,5-dimethoxy-phenyl)-1-prop-2-(E)-enyl	1,1-dimethylpropyl	1	0.450
	23	3-(3-pyridyl)-1-propyl	1,1-dimethylpropyl	1	0.0075
25	24	3-phenyl-1-propyl	cyclohexyl	1	0.082

TABLE II

30

	No.	m	n	Z	R ₂	R ₁	K _i nM
35	25	2	0	3-Phenylpropyl	3-(3-Pyridyl)propyl	Phenyl	56
	26	2	0	3-Phenylpropyl	3-(2-Pyridyl)	Phenyl	50

5	27	2	0	3-Phenylpropyl	propyl 2-(4-Methoxy-phenyl)ethyl	Phenyl	270
	28	2	0	3-Phenylpropyl	3-Phenylpropyl	Phenyl	---
	29	2	0	3-Phenylpropyl	3-Phenylpropyl	3,4,5- Trimethoxyphenyl	1.0
	30	2	0	3-Phenylpropyl	2-(3-Pyridyl)	3,4,5- Trimethoxyphenyl	3.0
	31	2	0	3-Phenylpropyl	3-(2-Pyridyl)	3,4,5- Trimethoxyphenyl	1.0
10	32	2	0	3-Phenylpropyl	3-(4-Methoxyphenyl)propyl	3,4,5- Trimethoxyphenyl	3.0
	33	2	0	3-Phenylpropyl	3-(3-Pyridyl)	3-Iso- propoxyphenyl	2.0
	34	1	1	3-pyridyl	3-phenyl	1,1-dimethylpropyl	0.019

Chick Dorsal Root Ganglion Cultures and Neurite Outgrowth

20 Dorsal root ganglia were dissected from chick embryos of ten day gestation.

Whole ganglion explants were cultured on thin layer Matrigel-coated 12 well plates with Liebovitz L15 plus high glucose media supplemented with 2mM glutamine and 10% fetal calf serum, and also containing 10 μ M cytosine β -D arabinofuranoside (Ara C) at 37°C in an environment containing 5% CO₂. Twenty-four hours later, the DRGs were

25 treated with various concentrations of nerve growth factor (NGF), immunophilin ligands or combinations of NGF plus drugs. Forty-eight hours after drug treatment, the ganglia were visualized under phase contrast or Hoffman Modulation contrast with a Zeiss Axiovert inverted microscope. Photomicrographs of the explants were made, and neurite outgrowth was quantitated. Neurites longer than the DRG diameter were

30 counted as positive, with total number of neurites quantitated per each experimental condition. Three to four DRGs were cultured per well, and each treatment was performed in duplicate.

The data for the drug alone (e.g., immunophilin ligand) experiments are presented in Table III.

TABLE III


Neurite Outgrowth in Chick DRG

Compound	ED ₅₀ , nM Neurite Outgrowth in DRG Cultures
1	25-100
2	10-20
3	0.500
4	25-100
5	25-100
6	10-20
7	>10,000
8	>10,000
9	>10,000
10	>10,000
11	1000
12	0.031
13	0.180
14	1-5
15	0.055
16	0.030
17	0.053
18	105
19	80
20	190
21	85
22	0.8
23	0.05
24	0.13
25	1-5
26	0.063
27	10-20
28	0.0044
29	0.61
30	0.95
31	25
32	0.50
33	0.30
34	0.07

EXAMPLES

The inventive compounds may be prepared by a variety of synthetic sequences that utilize established chemical transformations. The general pathway to the present compounds is described in Scheme 1. N-glyoxylproline derivatives may be prepared by reacting L-proline methyl ester with methyl oxalyl chloride as shown in Scheme I. The resulting oxamates may be reacted with a variety of carbon nucleophiles to obtain intermediates compounds. These intermediates are then reacted with a variety of alcohols, amides, or protected amino acid residues to obtain the propyl esters, ketones, acids, and amides of the invention.

Scheme I



10

EXAMPLE 1

Synthesis of methyl (2S)-1-(1,2-dioxo-2-methoxyethyl)-2-pyrrolidinecarboxylate.

A solution of L-proline methyl ester hydrochloride (3.08 g; 18.60 mmol) in dry methylene chloride was cooled to 0°C and treated with triethylamine (3.92 g; 38.74

mmol; 2.1 eq). After stirring the formed slurry under a nitrogen atmosphere for 15 min, a solution of methyl oxalyl chloride (3.20 g; 26.12 mmol) in methylene chloride (45 mL) was added dropwise. The resulting mixture was stirred at 0°C for 1.5 hr. After filtering to remove solids, the organic phase was washed with water, dried over MgSO₄ and concentrated. The crude residue was purified on a silica gel column, eluting with 50% ethyl acetate in hexane, to obtain 3.52 g (88%) of the product as a reddish oil. Mixture of cis-trans amide rotamers; data for trans rotamer given. ¹H NMR (CDCl₃): d 1.93 (dm, 2H); 2.17 (m, 2H); 3.62 (m, 2H); 3.71 (s, 3H); 3.79, 3.84 (s, 3H total); 4.86 (dd, 1H, J = 8.4, 3.3).

10

EXAMPLE 2

General procedure for the synthesis of pyrrolidinyl alkyl oxamates. Exemplified for methyl (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinecarboxylate.


A solution of methyl (2S)-1-(1,2-dioxo-2-methoxyethyl)-2-pyrrolidinecarboxylate (2.35 g; 10.90 mmol) in 30 mL of tetrahydrofuran (THF) was cooled to -78°C and treated with 14.2 mL of a 1.0M solution of 1,1-dimethylpropylmagnesium chloride in THF. After stirring the resulting homogenous mixture at -78°C for three hours, the mixture was poured into saturated ammonium chloride (100 mL) and extracted into ethyl acetate. The organic phase was washed with water, dried, and concentrated, and the crude material obtained upon removal of the solvent was purified on a silica gel column, eluting with 25% ethyl acetate in hexane, to obtain 2.10 g (75%) of the oxamate as a colorless oil. ¹H NMR (CDCl₃): d 0.88 (t, 3H); 1.22, 1.26 (s, 3H each); 1.75 (dm, 2H); 1.87-2.10 (m, 3H); 2.23 (m, 1H); 3.54 (m, 2H); 3.76 (s, 3H); 4.52 (dm, 1H, J=8.4, 3.4).

EXAMPLE 3

General procedure for the preparation of pyrrolidine carboxylic acids.

Exemplified for (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinecarboxylic acid.

A mixture of methyl (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinecarboxylate (2.10 g; 8.23 mmol), 1 N LiOH (15 mL), and methanol (50 mL) was stirred at 0°C for 30 min and at room temperature overnight. The mixture was acidified to pH 1 with 1 N HCl, diluted with water, and extracted into 100 mL of methylene chloride. The organic extract was washed with brine and concentrated to deliver 1.73 g (87%) of snow-white solid which did not require further purification. ¹H NMR (CDCl₃): δ 0.87 (t, 3H); 1.22, 1.25 (s, 3H each); 1.77 (dm, 2H); 2.02 (m, 2H); 2.17 (m, 1H); 2.25 (m, 1H); 3.53 (dd, 2H, J = 10.4, 7.3); 4.55 (dd, 1H, J = 8.6, 4.1).


EXAMPLE 4

Benzyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate.

A mixture of (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidine-carboxylic acid (500 mg; 2.07 mmol), benzyl alcohol (335 mg; 3.10 mmol), dicyclohexylcarbodiimide (683 mg; 3.31 mmol), 4-dimethylaminopyridine (84 mg; 0.69 mmol) and camphorsulphonic acid (160 mg; 0.69 mmol) in methylene chloride (30 mL) was stirred overnight under a nitrogen atmosphere. The reaction mixture was filtered through Celite


to remove solids and concentrated *in vacuo*, and the crude material was purified on a flash column (25% ethyl acetate in hexane) to obtain 680 mg of the product as a colorless oil. ^1H NMR (CDCl_3 ; 300 MHz) δ 0.85 (t, 3H); 1.19 (s, 3H); 1.22 (s, 3H); 1.61-2.25 (m, 6H); 3.46-3.56 (m, 2H); 4.58 (dm, 1H); 5.18 (d, 2H, 7.35 (br, 5H).

5


EXAMPLE 5

2-Phenyl-1-ethyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate.

A mixture of (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidine-carboxylic acid (570 mg; 2.36 mmol), phenethyl alcohol (432 mg; 3.54 mmol), dicyclohexylcarbodiimide (780 mg; 3.78 mmol), 4-dimethylaminopyridine (98 mg; 0.79 mmol) and camphorsulphonic acid (183 mg; 0.79 mmol) in methylene chloride (30 mL) was stirred overnight under a nitrogen atmosphere. The reaction mixture was filtered through Celite to remove solids and concentrated *in vacuo*, and the crude material was purified on a flash column (25% ethyl acetate in hexane) to obtain 600 mg of the product as a colorless oil. ^1H NMR (CDCl_3 ; 300 MHz) δ 0.87 (t, 3H); 1.21 (s, 3H); 1.25 (s, 3H); 1.64-1.94 (m, 6H); 2.17 (m, 1H); 2.97 (m, 2H); 3.49 (m, 2H); 4.36 (m, 2H); 4.51 (m, 1H); 7.22-7.33 (m, 5H).

EXAMPLE 6

General procedure for the synthesis of prolyl esters. Exemplified for 3-phenyl-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate. A mixture of (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidine-carboxylic acid (600 mg; 2.49 mmol), 3-phenyl-1-propanol (508 mg; 3.73 mmol), dicyclohexylcarbodiimide (822 mg; 3.98 mmol), camphorsulphonic acid (190 mg; 0.8 mmol) and 4-dimethylaminopyridine (100 mg; 0.8 mmol) in methylene chloride (20 mL) was stirred overnight under a nitrogen atmosphere. The reaction mixture was filtered through Celite to remove solids and concentrated in vacuo, and the crude material was purified on a flash column (25% ethyl acetate in hexane) to obtain 720 mg (80%) of Example 1 as a colorless oil. ¹H NMR (CDCl₃; 300 MHz): δ 0.84 (t, 3H); 1.19 (s, 3H); 1.23 (s, 3H); 1.70 (dm, 2H); 1.98 (m, 5H); 2.22 (m, 1H); 2.64 (m, 2H); 3.47 (m, 2H); 4.14 (m, 2H); 4.51 (d, 1H); 7.16 (m, 3H); 7.26 (m, 2H). Example 6 is compound 17 in Tables I and III.


EXAMPLE 7

3-phenyl-1-prop-2-(E)-enyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-

- 5 pyrrolidinecarboxylate, 80%, ^1H NMR (CDCl_3 ; 360 Mhz): δ 0.86 (t, 3H); 1.21 (s, 3H); 1.25 (s, 3H); 1.54-2.10 (m, 5H); 2.10-2.37 (m, 1H); 3.52-3.55 (m, 2H); 4.56 (dd, 1H, J = 3.8, 8.9); 4.78-4.83 (m, 2H); 6.27 (m, 1H); 6.67 (dd, 1H, J = 15.9); 7.13-7.50 (m, 5H).

This compound was prepared by the method of Example 3 from (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidine-carboxylic acid. Example 7 is compound 18 in Tables I

10 and III.


EXAMPLE 8

3-(3,4,5-trimethoxyphenyl)-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-

- 15 pyrrolidine- carboxylate, 61%, ^1H NMR (CDCl_3 ; 300 MHz): δ 0.84 (t, 3H); 1.15 (s, 3H); 1.24 (s, 3H); 1.71 (dm, 2H); 1.98 (m, 5H); 2.24 (m, 1H); 2.63 (m, 2H); 3.51 (t, 2H); 3.79 (s, 3H); 3.83 (s, 3H); 4.14 (m, 2H); 4.52 (m, 1H); 6.36 (s, 2H). This compound was

prepared by the method of Example 3 from (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidine-carboxylic acid. Example 8 is compound 19 in Tables I and III.

EXAMPLE 9




5

3-(3,4,5-trimethoxyphenyl)-1-prop-2-(E)-enyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidine carboxylate, 66%, ^1H NMR (CDCl_3 ; 360 MHz): δ 0.85 (t, 3H); 1.22 (s, 3H); 1.25 (s, 3H); 1.50-2.11 (m, 5H); 2.11-2.40 (m, 1H); 3.55 (m, 2H); 3.85 (s, 3H); 3.88 (s, 6H); 4.56 (dd, 1H); 4.81 (m, 2H); 6.22 (m, 1H); 6.58 (d, 1H, $J = 16$); 6.63 (s, 2H). This compound was prepared by the method of Example 3 from (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidine-carboxylic acid. Example 9 is compound 20 in Tables I and III.

10

EXAMPLE 10




15


3-(4,5-Dichlorophenyl)-1-prop-2-(E)-enyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-

pyrrolidine carboxylate, 70%, ^1H NMR (CDCl_3 ; 360 MHz): δ 0.85 (t, 3H); 1.21 (s, 3H); 1.25 (s, 3H); 1.51-1.87 (m, 2H); 1.87-2.39 (m, 4H); 3.51-3.57 (m, 2H); 4.50-4.61 (dd, 1H, $J=3.4, 8.6$); 4.80 (d, 2H, $J=6.0$); 6.20-6.34 (m, 1H); 6.50-6.66 (d, 1H, $J = 16$); 7.13-7.24 (dd, 1H, $J=1.8, 8.3$); 7.39 (d, 1H, $J=8.3$); 7.47 (s, 1H). This compound was prepared by the method of Example 3 from (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidine-carboxylic acid. Example 10 is compound 21 in Tables I and III.

EXAMPLE 11




(1R)-1,3-Diphenyl-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate, 90%, ^1H NMR (CDCl_3 ; 360 MHz): δ 0.85 (t, 3H); 1.20 (s, 3H); 1.23 (s, 3H); 1.49-2.39 (m, 7H); 2.46-2.86 (m, 2H); 3.25-3.80 (m, 2H); 4.42-4.82 (m, 1H); 5.82 (td, 1H, $J = 1.8, 6.7$); 7.05-7.21 (m, 3H); 7.21-7.46 (m, 7H). This compound was prepared by the method of Example 3 from (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidine-carboxylic acid.

Example 12

4-Phenyl-1-butyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate.


- 5 ^1H NMR (CDCl_3 ; 300 MHz): δ 0.84 (t, 3H); 1.22 (s, 3H); 1.25 (s, 3H); 1.64-2.01 (m, 9H); 2.23 (m, 1H); 2.64 (m, 2H); 3.48-3.53 (m, 2H); 4.17 (m, 2H); 4.52 (m, 1H); 7.18 (m, 3H); 7.27 (m, 2H).

Example 13

10 5-Phenyl-1-pentyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate.


- ^1H NMR (CDCl_3 ; 300 MHz): δ 0.87 (t, 3H); 1.22 (s, 3H); 1.25 (s, 3H); 1.39 (m, 2H); 1.63-1.99 (m, 9H); 2.22 (m, 1H); 2.64 (m, 2H); 3.46-3.54 (m, 2H); 4.14 (m, 2H); 4.50 (m, 1H); 7.16 (m, 3H); 7.26 (m, 2H).

- 15 Examples 12 and 13 were prepared according to the synthetic procedure outlined for Examples 1-3, except that the requisite phenyl alcohols in the reaction mixture were 4-phenylbutan-1-ol and 5-phenylpentan-1-ol, respectively.

Example 14


1,1-Diphenylmethyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate.

^1H NMR (CDCl_3 ; 300 MHz): δ 0.84 (t, 3H); 1.17 (s, 3H); 1.19 (s, 3H); 1.54-2.25 (m, 6H); 3.50 (m, 2H); 4.67 (m, 1H); 5.86 (s, 1H); 7.28-7.39 (m, 10H).

Example 15

1,3-diphenyl-2-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate.


^1H NMR (CDCl_3 ; 300 MHz): δ 0.87 (t, 3H); 1.20 (s, 3H); 1.24 (s, 3H); 1.25-2.02 (m, 6H); 2.74, 2.84 (m, 4H total); 3.53 (m, 2H); 4.04 (m, 1H); 4.42 (m, 1H); 7.22 (m, 6H); 7.30 (m, 4H).

Example 16

1,5-diphenyl-3-pentyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-

5 pyrrolidinecarboxylate.

^1H NMR (CDCl_3 ; 300 MHz): δ 0.87 (t, 3H); 1.23 (s, 3H); 1.27 (s, 3H); 1.61-2.06 (m, 9H); (m, 2H); 2.28 (m, 1H); 2.57-2.74 (m, 4H); 3.52-3.56 (m, 2H); 4.49-4.59 (m, 1H); 5.02 (m, 1H); 7.14-7.30 (m, 10H).

Example 17

1,7-diphenyl-4-heptyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-


pyrrolidinecarboxylate.

^1H NMR (CDCl_3 ; 300 MHz): δ 0.86 (t, 3H); 1.23 (s, 3H); 1.25 (s, 3H); 1.44-1.98 (m, 13H); 2.21 (m, 1H); 2.59 (m, 4H); 3.45-3.63 (m, 2H); 4.48-4.52 (dd, 1H, $J=2.7, 6.5$); 4.99 (m, 1H); 7.08-7.18 (m, 6H); 7.21-7.29 (m, 4H).

Examples 14-17 were prepared according to the synthetic procedure outlined for

Examples 1-3, except that the requisite diphenyl alcohols in the reaction mixture were 1,1-diphenylmethanol, 1,3-diphenylpropan-2-ol, 1,5-diphenylpentan-3-ol, and 1,7-diphenylheptan-4-ol, respectively.

Example 18




5

3-(2,5-Dimethoxyphenyl)-1-prop-2(*E*)-enyl (2*S*)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate.

¹H NMR (CDCl₃; 300 MHz): δ 0.87 (t, 3H); 1.22 (s, 3H); 1.26 (s, 3H); 1.67 (m, 2H); 1.78 (m, 1H); 2.07 (m, 2H); 2.26 (m, 1H); 3.52 (m, 2H); 3.78 (s, 3H); 3.80 (s, 3H); 4.54 (m, 1H); 4.81 (m, 2H); 6.29 (dt, 1H, *J*=15.9); 6.80 (s, 2H); 6.95 (d, 1H, *J*=15.9); 6.98 (s, 1H). Example 18 is compound 22 in Tables I and III.

10

Example 19



15


3-Cyclohexyl-1-prop-2(*E*)-enyl (2*S*)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-

pyrrolidinecarboxylate.

^1H NMR (CDCl_3 ; 360 MHz): δ 0.86 (t, 3H); 1.13-1.40 (m +2 singlets, 9H total); 1.50-1.87 (m, 8H); 1.87-2.44 (m, 6H); 3.34-3.82 (m, 2H), 4.40-4.76 (m, 3H); 5.35-5.60 (m, 1H); 5.60-5.82 (dd, 1H, $J=6.5$, 16).

- 5 Examples 18 and 19 were prepared according to the synthetic procedure outlined for Examples 1-3, except that the requisite trans-allylic alcohols in the reaction mixture were 3-(2,5-Dichlorophenyl)-1-prop-2(*E*)-enol and 3-Cyclohexyl-1-prop-2(*E*)-enol, respectively.


Example 20



10

(3,4,5-Trimethoxy)benzyl (2*S*)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate.


- 15 ^1H NMR (CDCl_3 ; 300 MHz): δ 0.85 (t, 3H); 1.20 (s, 3H); 1.22 (s, 3H); 1.58-1.81 (m, 2H); 1.82-2.27 (m, 4H); 3.52 (m, 2H); 3.84 *s, 3H); 3.87 (s, 6H); 4.55 (m, 1H); 5.13 (s, 2H); 6.59 (s, 2H).

Example 21

5 2-(3,4,5-Trimethoxyphenyl)-1-ethyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate

^1H NMR (CDCl_3 ; 300 MHz): δ 0.84 (t, 3H); 1.15 (s, 3H); 1.24 (s, 3H); 1.71 (dm, 2H); 1.98 (m, 5H); 2.24 (m, 1H); 2.63 (m, 2H); 3.51 (t, 2H); 3.79 (s, 3H); 3.83 (s, 3H); 4.14 (m, 2H); 4.52 (m, 1H); 6.36 (s, 2H).

10

Example 22


3-(2,5-Dimethoxyphenyl)-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate.

15 ^1H NMR (CDCl_3 ; 300 MHz): δ 0.87 (t, 3H); 1.22 (s, 3H); 1.26 (s, 3H); 1.69 (m, 2H); 1.96 (m, 5H); 2.24 (m, 1H); 2.68 (m, 2H); 3.55 (m, 2H); 3.75 (s, 3H); 3.77 (s, 3H);

4.17 (m, 2H); 4.53 (d, 1H); 6.72 (m, 3H).

Examples 20, 21, and 22 were prepared according to the synthetic procedure outlined for Examples 1-3, except that the requisite di- or trimethoxyphenyl-substituted alcohols in the reaction mixture were (3,4,5-Trimethoxy)benzyl alcohol, 2-(3,4,5-Trimethoxyphenyl)-1-ethanol, and 3-(2,5-Dimethoxyphenyl)-propan-1-ol, respectively.

Example 23




(1S)-1,3-Diphenyl-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-

pyrrolidinecarboxylate.

^1H NMR (CDCl_3 ; 360 MHz): δ 0.87 (t, 3H); 1.20 (s, 3H); 1.24 (s, 3H); 1.62-2.32 (m, 8H); 2.62-2.75 (m, 2H); 3.43-3.60 (m, 2H); 4.58-4.73 (m, 1H); 5.76 (td, 1H, $J=1.8$, 6.7); 7.19 (m, 3H); 7.24-7.35 (m, 7H).


Example 23 was prepared according to the synthetic procedure outlined for Examples 1-3, except that the requisite optically active 1-substituted alkanol in the reaction mixture was (1S)-1,3-Diphenylpropan-1-ol.

Example 24

3-(3-Pyridyl)-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate.

^1H NMR (CDCl_3 ; 360 MHz): δ 0.85 (t, 3H); 1.23, 1.26 (s, 3H each); 1.69-1.90 (m, 3H); 1.95-2.01 (m, 4H); 2.20 (m, 1H); 2.72 (t, 2H); 3.53 (m, 2H); 4.18 (m, 2H); 4.52 (m, 1H); 7.22 (m, 1H); 7.53 (dd, 1H); 8.45 (m, 2H).

Example 24 was prepared according to the synthetic procedure outlined for Examples 1-3, except that the requisite alcohol in the reaction mixture was 3-(3-Pyridyl)propan-1-ol. Example 24 is compound 23 in Tables I and III.

Example 25

3-Phenyl-1-propyl (2S)-1-(cyclohexylglyoxyl)-2-pyrrolidinecarboxylate.


^1H NMR (CDCl_3 ; 300 MHz): δ 1.09-1.33 (m, 5H); 1.62-2.33 (m, 12H); 2.69 (t, 2H, $J=7.5$); 3.15 (dm, 1H); 3.68 (m, 2H); 4.16 (m, 2H); 4.53, 4.84 (d, 1H total); 7.19 (m, 3H);

7.29 (m, 2H). Example 25 is compound 24 in Tables I and III.

EXAMPLE 26

The requisite substituted alcohols may be prepared by a number of methods known to those skilled in the art of organic synthesis. As described in Scheme II, alkyl or aryl aldehydes may be homologated to phenyl propanols by reaction with methyl (triphenylphosphoranylidene) acetate to provide a variety of *trans*-cinnamates; these latter may be reduced to the saturated alcohols by reaction with excess lithium aluminum hydride, or sequentially by reduction of the double bond by catalytic hydrogenation and reduction of the saturated ester by appropriate reducing agents. Alternatively, the *trans*-cinnamates may be reduced to (E)-allylic alcohols by the use of diisobutylaluminum hydride.

Scheme II



Longer chain alcohols may be prepared by homologation of benzylic and higher aldehydes. Alternatively, these aldehydes may be prepared by conversion of the corresponding phenylacetic and higher acids, and phenethyl and higher alcohols.

EXAMPLE 27

General procedure for the synthesis of acrylic esters, exemplified for methyl (3,3,5-trimethoxy)-*trans*-cinnamate:

A solution of 3,4,5-trimethoxybenzaldehyde (5.0 g; 25.48 mmol) and methyl (triphenyl- phosphoranylidene)acetate (10.0 g; 29.91 mmol) in tetrahydrofuran (250 mL) was refluxed overnight. After cooling, the reaction mixture was diluted with 200 mL of ethyl acetate and washed with 2 x 200 mL of water, dried, and concentrated in vacuo.

- 5 The crude residue was chromatographed on a silica gel column, eluting with 25% ethyl acetate in hexane, to obtain 5.63 g (88%) of the cinnamate as a white crystalline solid, ^1H NMR (300 Mhz; CDCl_3): δ 3.78 (s, 3H); 3.85 (s, 6H); 6.32 (d, 1H, $J = 16$); 6.72 (s, 2H); 7.59 (d, 1H, $J = 16$).

EXAMPLE 28

- 10 Methyl (4,5-dichloro)-trans-cinnamate, 80%, ^1H NMR (300 Mhz; CDCl_3): δ 3.79 (s, 3H); 6.40 (d, 1H, $J=16.8$); 7.32 (dd, 1H, $J = 1.5, 8.1$); 7.44 (d, 1H, $J=8.1$); 7.56 (d, 1H, $J = 16$); 7.58 (s, 1H). This compound was prepared by the method of Example 27 from 3,4,5-trimethoxybenzaldehyde.

EXAMPLE 29

- 15 Methyl (4,5-methylenedioxy)-trans-cinnamate, 74%, ^1H NMR (360 Mhz; CDCl_3): δ 3.79 (s, 3H); 6.01 (s, 2H); 6.26 (d, 1H, $J=16$); 6.81 (d, 1H, $J=7.9$); 7.00 (d, 1H, $J=8.2$); 7.03 (s, 1H); 7.60 (d, 1H, $J=16$). This compound was prepared by the method of Example 27 from 3,4,5-trimethoxybenzaldehyde.

EXAMPLE 30

- 20 Methyl (2-cyclohexyl)-(E)-acrylate, 80%, ^1H NMR (360 Mhz; CDCl_3): δ 1.12-1.43 (m, 5H); 1.52-1.87 (m, 5H); 2.12 (m, 1H); 2.12 (m, 1H); 3.71 (s, 3H); 5.77 (dd, 1H, $J=1.2, 15.8$); 6.92 (dd, 1H, $j=6.8, 15.8$). This compound was prepared by the method of Example 27 from 3,4,5-trimethoxybenzaldehyde.

EXAMPLE 31

General procedure for the synthesis of saturated alcohols from acrylic esters.

Exemplified for (3,4,5-trimethoxy)phenylpropanol.

A solution of methyl (3,3,5-trimethoxy)-*trans*-cinnamate (1.81 g; 7.17 mmol) in
 5 tetrahydrofuran (30 mL) was added in a dropwise manner to a solution of lithium
 aluminum hydride (14 mmol) in THF (35 mL), with stirring and under an argon
 atmosphere. After the addition was complete, the mixture was heated to 75°C for 4
 hours. After cooling, it was quenched by the careful addition of 15 mL of 2N NaOH
 followed by 50 mL of water. The resulting mixture was filtered through Celite to remove
 10 solids, and the filter cake was washed with ethyl acetate. The combined organic
 fractions were washed with water, dried, concentrated in vacuo, and purified on a silica
 gel column, eluting with ethyl acetate to obtain 0.86 g (53%) of the alcohol as a clear oil,
¹H NMR (300 Mhz; CDCl₃): δ 1.23 (br, 1H); 1.87 (m, 2H); 2.61 (t, 2H, J = 7.1); 3.66 (t,
 2H); 3.80 (s, 3H); 3.83 (s, 6H); 6.40 (s, 2H).

EXAMPLE 32

General procedure for the synthesis of *trans*-allylic alcohols from acrylic esters.

Exemplified for (3,4,5-trimethoxy)phenylprop-2-(E)-enol.

A solution of methyl (3,3,5-trimethoxy)-*trans*-cinnamate (1.35 g; 5.35 mmol) in
 toluene (25 mL) was cooled to -10°C and treated with a solution of diisobutylaluminum
 20 hydride in toluene (11.25 mL of a 1.0 M solution; 11.25 mmol). The reaction mixture was
 stirred for 3 hrs at 0°C and then quenched with 3 mL of methanol followed by 1 N HCl
 until the pH was 1. The reaction mixture was extracted into ethyl acetate and the
 organic phase was washed with water, dried and concentrated. Purification on a silica

gel column eluting with 25% ethyl acetate in hexane furnished 0.96 g (80%) of a thick oil, ^1H NMR (360 Mhz; CDCl_3): δ 3.85 (s, 3H); 3.87 (s, 6H); 4.32 (d, 2H, $J = 5.6$); 6.29 (dt, 1H, $J = 15.8, 5.7$), 6.54 (d, 1H, $J = 15.8$); 6.61 (s, 2H).

EXAMPLE 33

5 (4,5-dichloro)phenylprop-2-(E)-enol, 89%, ^1H NMR (360 Mhz; CDCl_3): δ 1.55 (s, 3H); 4.34 (d, 2H, $J=4.4$); 6.36 (dt, 1H, $J=15.9, 5.3$); 6.54 (d, 1H, $J=15.9$); 7.20 (dd, 1H, $J=8.3, 1.7$); 7.38 (d, 1H, $J=8.3$); 7.45 (d, 1H, $J=1.6$). This compound was prepared by the method of Example 32 from 3,4,5-(trimethoxy)-trans-cinnamate.

EXAMPLE 34

10 (4,5-methylenedioxy)phenylprop-2-(E)-enol, 80%, ^1H NMR (360 Mhz; CDCl_3): δ 1.59 (br, 1H); 4.29 (br, 2H); 5.96 (s, 2H); 6.20 (dt, 1H, $J=15.8, 5.9$); 6.52 (d, 1H, $J=15.8$); 6.76 (d, 1H, $J=8.0$); 6.82 (dd, 1H, $J=8.0, 1.2$); 6.93 (d, 1H, $J=1.2$). This compound was prepared by the method of Example 32 from 3,4,5-(trimethoxy)-trans-cinnamate.

EXAMPLE 35


15 Phenylprop-2-(E)-enol, 85%, ^1H NMR (360 Mhz; CDCl_3): δ 1.72 (br, 1H); 4.31 (d, 2H, $J=5.7$); 6.36 (dt, 1H, $J=15.9, 5.7$); 6.61 (d, 1H, $J=15.9$); 7.02-7.55 (m, 5H). This compound was prepared by the method of Example 32 from 3,4,5-(trimethoxy)-trans-cinnamate.

EXAMPLE 36

20 Alcohols containing a substituent at the 1-position of the side chain may be conveniently prepared by addition of appropriate nucleophiles to aldehydes, as described in Scheme III. In cases where optically active substituted alcohols are desired, the racemic alcohols may be oxidized to prochiral ketones and subjected to

asymmetric reduction by one of several methods well known to those skilled in the art.

Scheme III



EXAMPLE 37

5 General procedure for the preparation of 1-substituted alkanols, exemplified for the synthesis of 1,3-diphenylpropanol.


A solution of 2-(bromoethyl)benzene (17.45 g; 94.3 mmol) in 50 mL of dry diethyl ether was added dropwise, under a nitrogen atmosphere, to a stirred slurry of magnesium turnings (2.50 g; 102.8 mmol) in 50 mL of ether. The mixture was initially
 10 heated with a heat gun until reflux had become self-sustaining. After the addition was complete, the mixture was heated externally for 30 min to maintain reflux. A solution of 10.01 g (94.3 mmol) of benzaldehyde in 20 mL of ether was then added dropwise, and reflux was continued for 30 min. After cooling, the reaction mixture was poured into 150 mL of saturated ammonium chloride and extracted into ethyl acetate. The crude
 15 material obtained upon removal of the solvent was purified on a flash column, eluting with 5% ethyl acetate/hexane to 20% ethyl acetate, to obtain 13.73 g (69%) of the alkanol as a light yellow oil, ^1H NMR (360 Mhz; CDCl_3): δ 1.93-2.30 (m, 3H); 2.70-2.90 (m, 2H); 4.72 (br, 1H); 7.19-7.27 (m, 3H); 7.27-7.36 (m, 3H); 7.36-7.47 (m, 4H).

EXAMPLE 38


General procedure for conversion of racemic 1-substituted alkanols to optically active 1-substituted alkanols via prochiral ketones. Exemplified for (1R)-1,3-diphenyl-1-propanol.

5 A solution of racemic 1,3-diphenyl-1-propanol (1.26 g; 5.94 mmol) was dissolved in 10 mL of acetone, and Jones reagent was added until persistence of the orange color. After stirring for 30 min, the reaction was quenched by adding 2 mL of 2-propanol. The solvent was decanted away from the precipitated solids, which were washed with ethyl acetate. The combined organic fractions were washed with 2x20 mL
10 of water, dried and concentrated. The crude product was filtered through a plug of silica gel, eluting with 25% ethyl acetate/hexane, to obtain 1.07 g (86%) of 1,3-diphenylpropanone as a white crystalline solid, ^1H NMR (360 Mhz; CDCl_3): δ 3.09 (t, 2H, $J=8.1$); 3.33 (t, 2H, $J=8.1$); 7.29 (m, 5H); 7.49 (m, 3H); 7.98 (m, 2H).

 A solution of 1,3-diphenylpropanone (1.07 g; 5.09 mmol) in tetrahydrofuran (10
15 mL) was cooled to -23°C and treated with an asymmetric reducing agent, (+)-B-chlorodiisopinocampheyl-borane (1.80 g; 5.60 mmol) in 20 mL THF, and the resulting solution was allowed to stand overnight at -23°C . After evaporating to dryness, the residue was treated with ether (65 mL) and diethanolamine (1.0 g) and stirred for 3 hrs. The mixture was then filtered to remove solids and concentrated, and the residue was
20 purified using gradient elution (5% ethyl acetate/hexane to 10% ethyl acetate) on a silica gel column to obtain 660 mg (61%) of (1R)-1,3-diphenyl-1-propanol as a crystalline white solid, ^1H NMR (360 Mhz; CDCl_3): δ 1.95-2.15 (m, 3H); 2.59-2.78 (m, 2H); 4.65 (dd, 1H, $J=5.4, 7.8$); 7.14-7.35 (m, 10H).

Scheme IVExample 39


5 Synthesis of ethyl 1-(1,2-dioxo-2-methoxyethyl)-2-piperidinecarboxylate.



A solution of ethyl pipercolinate (1.00 g; 5.57 mmol) in dry methylene chloride (15 mL) was cooled to 0°C and treated with triethylamine (1.24 g; 12.25 mmol; 2.1 eq). After stirring the formed slurry under a nitrogen atmosphere for 15 min, a solution of methyl oxalyl chloride (0.96 g; 6.13 mmol) in methylene chloride (15 mL) was added dropwise. The resulting mixture was stirred at 0°C for 1.5 hr. After filtering to remove solids, the organic phase was washed with water, dried over MgSO_4 and concentrated. The crude residue was purified on a silica gel column, eluting with 50% ethyl acetate in hexane, to

obtain 1.21 g (95%) of the product as a reddish oil. Mixture of cis-trans amide rotamers; data for trans rotamer given. ^1H NMR (300 MHz, CDCl_3): δ 1.25 (t, 3H); 1.30-1.75 (m, 5H); 2.33 (m, 1H); 3.42 (dt, 1H); 3.57 (br d, 1H); 3.85 (s, 3H); 4.29 (dd, 2H); 5.23 (d, 1H).

5


Example 40

Synthesis of ethyl 1-(1,2-dioxo-3,3-dimethylpentyl)-2-piperidinecarboxylate.


10

A solution of ethyl 1-(1,2-dioxo-2-methoxyethyl)-2-piperidinecarboxylate (1.43 g; 5.88 mmol) in 20 mL of tetrahydrofuran (THF) was cooled to -78°C and treated with 8 mL of a 1.0 M solution of 1,1-dimethylpropylmagnesium chloride in THF. After stirring the resulting homogeneous mixture at -78°C for three hours, the mixture was poured into saturated ammonium chloride (30 mL) and extracted into ethyl acetate. The organic phase was washed with water, dried, and concentrated, and the crude material obtained upon removal of the solvent was purified on a silica gel column, eluting with 25% ethyl acetate in hexane, to obtain 1.35 g (76%) of the oxamate as a colorless oil. ^1H NMR (300 MHz, CDCl_3): δ 0.91 (t, 3H); 1.20, 1.25 (s, 3H each); 1.30 (t, 3H); 1.35-1.80 (m, 7H); 2.35 (br d, 1H); 3.20 (td, 1H); 3.41 (br d, 1H); 4.20 (q, 2H); 5.22 (d, 1H).

20

Example 41**Synthesis of 1-(1,2-dioxo-3,3-dimethylpentyl)-2-piperidinecarboxylic acid**

- 5 A mixture of ethyl 1-(1,2-dioxo-3,3-dimethylpentyl)-2-piperidinecarboxylate (0.69 g; 2.43 mmol), 1 N LiOH (5 mL), and methanol (20 mL) was stirred at 0°C for 30 min and at room temperature overnight. The mixture was acidified to pH 1 with 1 N HCl, diluted with water, and extracted into 50 mL of methylene chloride. The organic extract was washed with brine and concentrated to deliver 0.61 g (98%) of snow-white solid
- 10 which did not require further purification.

Example 42**Synthesis of 3-Phenyl-1-propyl 1-(1,2-dioxo-3,3-dimethylpentyl)-2-**


- 15 piperidinecarboxylate

- A mixture of 1-(1,2-dioxo-3,3-dimethylpentyl)-2-piperidinecarboxylic acid (590 mg; 2.31 mmol), 3-phenylpropanol (520 mg; 3.71 mmol), dicyclohexylcarbodiimide (815 mg; 3.95 mmol), camphorsulphonic acid (180 mg; 0.77 mmol) and 4-dimethyl aminopyridine (95 mg; 0.77 mmol) in methylene chloride (15 mL) was stirred overnight
- 20 under a nitrogen atmosphere. The reaction mixture was filtered through Celite to remove solids and concentrated in vacuo. The crude material was triturated with several

portions of ether, and the ether portions were filtered through Celite to remove solids and concentrated in vacuo. The concentrated filtrate was purified on a flash column (20% ethyl acetate in hexane) to obtain 800 mg (93%) of the product as an oil, ^1H NMR (CDCl_3 , 300 MHz): δ 0.85 (t, 3H); 1.23, 1.26 (s, 3H each); 1.63-1.94 (m, 9H); 2.32 (m, 1H); 2.69 (m, 2H); 3.21 (m, 1H); 3.35 (m, 1H); 4.17 (m, 2H); 5.24 (m, 1H); 7.14 (m, 3H); 7.7.23 (m, 2H).


The following compounds were prepared by this method:

Example 43



10 3-(3,4,5-Trimethoxyphenyl)-1-propyl 1-(3,3-dimethyl-1,2-dioxopentyl)-2-piperidinecarboxylate: ^1H NMR (CDCl_3 , 300 MHz): δ 0.80 (t, 3H); 1.18 (s, 6H); 1.67 (m, 7H); 1.94 (m, 2H); 2.29 (br d, 1H); 2.61 (t, 2H); 3.17 (td, 1H); 3.35 (d, 1H); 3.79 (s, 3H); 3.81 (s, 6H); 4.15 (m, 2H); 5.24 (d, 1H).


Example 44



15 3-(4,5-Dichlorophenyl)-1-prop-2-(E)-enyl 1-(3,3-dimethyl-1,2-dioxopentyl)-2-piperidinecarboxylate: ^1H NMR (CDCl_3 , 360 MHz): δ 0.89 (t, 3H); 1.18 (s, 3H); 1.24 (s, 3H); 1.57-1.89 (m, 7H); 2.38 (d, 1H); 3.20-3.28 (dt, 1H); 3.30-3.43 (dm, 1H); 4.81 (d,


2H); 5.31 (d, 1H); 6.16-6.36 (m, 1H); 6.48-6.68 (d, 1H); 7.20 (d, 1H); 7.39 (d, 1H); 7.47 (s, 1H).

Example 45




5 3-(3,4,5-Trimethoxyphenyl)-1-prop-2-(E)-enyl 1-(3,3-dimethyl-1,2-dioxopentyl)-2-piperidinecarboxylate: ^1H NMR (CDCl_3 , 360 MHz): δ 0.89 (t, 3H); 1.21 (s, 3H); 1.24 (s, 3H); 1.41-1.85 (m, 7H); 2.35 (d, 1H); 3.25 (t, 1H); 3.39 (m, 1H); 3.86 (s, 3H); 3.89 (s, 6H); 4.81 (m, 2H); 5.33 (d, 1H); 6.21 (m, 1H); 6.61 (d, 1H); 6.63 (s, 2H).

Example 46




10 3-Phenyl-1-prop-2-(E)-enyl 1-(3,3-dimethyl-1,2-dioxopentyl)-2-piperidinecarboxylate: ^1H NMR (CDCl_3 , 360 MHz): δ 0.88 (t, 3H); 1.20 (s, 3H); 1.24 (s, 3H); 1.25-1.77 (m, 6H); 1.86-2.06 (m, 1H); 2.30-2.40 (m, 1H); 3.24 (t, 1H); 3.41 (d, 1H); 4.82 (d, 1H); 5.31 (d, 1H); 6.25-6.29 (m, 1H); 6.68 (d, 1H); 7.26-7.54 (m, 5H).


15

Example 47

4-(4-Methoxyphenyl)butyl N-(phenylglyoxyl)-2-piperidinecarboxylate: ^1H NMR (CDCl₃, 300 MHz): δ 1.26-1.78 (m, 9H); 2.36 (d, 1H); 2.58 (m, 2H); 3.25 (m, 1H); 3.48 (dm, 1H); 3.78 (s, 3H); 4.24 (m, 2H); 5.40 (m, 1H); 6.82 (d, 2H); 7.09 (d, 2H); 7.64 (m, 2H); 7.66 (m, 1H); 8.02 (m, 2H). Anal. Calcd. for C₂₅H₂₉NO₅: C, 70.90; H, 6.90; N, 3.31. Found: C, 70.87; H, 6.92; N, 3.36. Example 47 is compound 3 in Tables I and III.

Example 48


1,7-Diphenylheptanyl N-(phenylglyoxyl)-2-piperidinecarboxylate: ^1H NMR (CDCl₃, 300 MHz): δ 1.21-1.85 (m, 14H); 2.48 (m, 4H); 3.22 (m, 1H); 3.44 (m, 1H); 5.09 (br, 1H); 5.38 (br, 1H); 7.06-8.04 (m, 15H). Anal. Calcd. for C₃₃H₃₇NO₄: C, 77.47; H, 7.29; N, 2.74. Found: C, 77.39; H, 7.32; N, 2.66. Example 48 is compound 20 in Tables I and III.

Example 49


3-Phenyl-1-propyl N-(phenylglyoxyl)-2-piperidinecarboxylate: ^1H NMR (CDCl₃,

300 MHz): δ 1.36-2.05 (m, 7H); 2.36 (d, 1H); 2.74 (m, 2H); 3.24 (t, 1H); 3.50 (t, 1H); 4.25 (m, 2H); 5.42 (m, 1H); 7.28 (m, 4H); 7.64 (m, 4H); 8.03 (m, 2H). Anal. Calcd. for $C_{23}H_{25}NO_4$: C, 72.80; H, 6.64; N, 3.69. Found: C, 72.74; H, 6.62; N, 3.62. Example 49 is compound 2 in Tables I and III.


5

Example 50

3-(3-Pyridyl)-1-propyl N-(phenylglyoxyl)-2-piperidinecarboxylate: 1H NMR ($CDCl_3$, 300 MHz): δ 1.26-2.08 (m, 7H); 2.35 (d, 1H); 2.75 (t, 2H); 3.29 (t, 1H); 3.49 (d, 1H); 4.27 (t, 2H); 5.42 (d, 1H); 7.23 (m, 1H); 7.52 (m, 3H); 7.63 (m, 1H); 8.03 (m, 2H); 8.48 (m, 2H). Anal. Calcd. for $C_{22}H_{24}N_2O_4 \cdot 0.25 H_2O$: C, 68.64; H, 6.42; N, 7.28. Found: C, 68.37; H, 6.41; N, 7.22.


Example 51

4-Phenyl-1-butyl N-(phenylglyoxyl)-2-piperidinecarboxylate: 1H NMR ($CDCl_3$, 300 MHz): δ 1.26-1.80 (m, 12H); 2.67 (m, 2H); 3.23 (t, 1H); 3.49 (t, 1H); 4.25 (m, 2H); 5.40 (m, 1H); 7.18 (m, 3H); 7.26 (m, 2H); 7.48 (m, 2H); 7.64 (m, 1H); 8.03 (m, 2H). Anal. Calcd. for $C_{24}H_{27}NO_4$: C, 73.26; H, 6.92; N, 3.56. Found: C, 73.19; H, 6.94; N, 3.64. Example 51 is compound 4 in Tables I and III.


Example 52

2-Phenyl-1-ethyl N-(phenylglyoxyl)-2-piperidinecarboxylate: ^1H NMR (CDCl_3 , 300 MHz): δ 1.23-1.75 (m, 5H); 2.21 (d, 1H); 3.09 (m, 3H); 3.41 (d, 1H); 4.48 (m, 2H); 5.38 (m, 1H); 7.27 (m, 5H); 7.53 (m, 2H); 7.65 (m, 1H); 8.01 (m, 2H). Anal. Calcd. for $\text{C}_{22}\text{H}_{23}\text{NO}_4 \cdot 0.25\text{H}_2\text{O}$: C, 71.43; H, 6.40; N, 3.79. Found: C, 71.60; H, 6.50; N, 4.12.

Example 52 is compound 5 in Tables I and III.

Example 53

Benzyl N-(phenylglyoxyl)-2-piperidinecarboxylate: ^1H NMR (CDCl_3 , 300 MHz): δ 1.38-1.81 (m, 5H); 2.41 (d, 1H); 3.22 (m, 1H); 3.48 (d, 1H); 5.26 (s, 2H); 5.47 (d, 1H); 7.42 (m, 7H); 7.61 (m, 1H); 7.97 (m, 2H). Anal. Calcd. for $\text{C}_{21}\text{H}_{21}\text{NO}_4 \cdot 0.25\text{H}_2\text{O}$: C, 71.78; H, 6.02; N, 3.99. Found: C, 71.90; H, 6.12; N, 4.01. Example 53 is compound 1 in Tables I and III.


Example 54

Benzyl N-(methoxyglyoxyl)-2-piperidinecarboxylate: ^1H NMR (CDCl_3 , 300 MHz): δ 1.26-1.77 (m, 5H); 2.32 (m, 1H); 3.33 (t, 1H); 3.54 (d, 1H); 3.88 (s, 3H); 5.23 (s, 2H);

5.45 (m, 1H); 7.36 (s, 5H). Anal. Calcd. for $C_{16}H_{19}NO_5$: C, 62.94; H, 6.27; N, 4.59.

Found: C, 62.80; H, 6.35; N, 4.53.

Scheme V




5

Example 55

Synthesis of (S)-Boc-pipecolyl-1,7-diphenyl-4-heptanyl ester:


A solution of (S)-boc-pipecolic acid (330 mg; 1.44 mmol) in CH_2Cl_2 (20 mL) was treated with 1,7-diphenyl-4-heptanol (350 mg; 1.30 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (280 mg; 1.44 mmol), and a catalytic amount of N,N,-dimethylaminopyridine. The reaction mixture was stirred overnight at room temperature, concentrated, and purified on a silica gel column eluting with 25% ethyl acetate in hexanes to provide 160 mg of product as a clear oil.

(S)-1,7-diphenyl-4-heptanylpipecolate: A solution of (S)-boc-pipecolyl-1,7-diphenyl-4-heptanyl ester (150 mg) in 10 mL of CH_2Cl_2 was treated with 3 mL of trifluoroacetic acid and stirred at room temperature for 2 hours. It was neutralized with aqueous potassium carbonate and the layers were separated. The organic phase was dried over $MgSO_4$ and concentrated to provide 70 mg of the free amine.

Example 56

1,7-Diphenyl-4-heptyl (S)-N-(3,4,5-trimethoxyphenylglyoxyl)pipecolate: A solution of (S)-1,7-diphenyl-4-heptylpipecolate (50 mg; 0.13 mmol) and 3,4,5-trimethoxybenzoyl- formic acid (45 mg; 0.2 mmol) was treated with 1-(3-dimethylaminopropyl)-3-ethyl- carbodiimide hydrochloride (40 mg; 0.2 mmol) and a catalytic amount of N,N,-dimethylaminopyridine. The reaction mixture was stirred overnight at room temperature, concentrated, and purified on a silica gel column eluting with 25% ethyl acetate in hexanes to provide 20 mg of product as a clear oil, ^1H NMR (300 MHz; CDCl_3): δ 1.31-1.92 (m, 13H); 2.35 (m, 1H); 2.66 (m, 4H); 3.29 (td, 1H); 3.94 (s, 9H); 5.08 (m, 1H); 5.41 (d, 1H); 7.19 (m, 6H); 7.28 (m, 4H); 7.42 (m, 2H). Example 56 is compound 21 in Tables I and III.

The following compounds were prepared by the method of Scheme V:


Example 57

15

3-(Phenoxybenzyl) (S)-N-(3,4,5-trimethoxyphenylglyoxyl)pipecolate: ^1H NMR (300 MHz; CDCl_3): δ 1.22-1.47 (m, 1H); 1.50-1.70 (m, 2H); 1.72-1.93 (m, 2H); 2.39 (d, 1H); 3.25 (td, 1H); 3.51 (d, 1H); 3.96 (s, 9H); 5.18 (m, 2H); 5.43 (d, 1H); 7.01 (m, 4H);

7.15 (m, 3H); 7.37 (m, 4H). Example 57 is compound 13 in Tables I and III.


Example 58




4-Phenylbutyl (S)-N-(3,4,5-trimethoxyphenylglyoxyl)pipecolate: ^1H NMR (300 MHz; CDCl_3): δ 1.32-1.88 (m, 9H); 2.35 (d, 1H); 2.63 (m, 2H); 3.25 (td, 1H); 3.48 (d, 1H); 3.93 (s, 9H); 4.18 (m, 2H); 5.35 (d, 1H); 7.17 (m, 3H); 7.23 (m, 2H); 7.36 (s, 2H).

Example 58 is compound 14 in Tables I and III.

Example 59




4-(4-Methoxyphenyl)butyl (S)-N-(3,4,5-trimethoxyphenylglyoxyl)pipecolate: ^1H NMR (300 MHz; CDCl_3): δ 1.21-1.92 (m, 9H); 2.37 (m, 1H); 2.62 (m, 2H); 3.25 (td, 1H); 3.49 (d, 1H); 3.78 (s, 3H); 3.93 (s, 9H); 4.15-4.23 (m, 2H); 5.38 (m, 1H); 6.39 (m, 2H); 7.07 (m, 1H); 7.36 (m, 1H). Example 59 is compound 16 in Tables I and III.

Example 60


1-Phenyl-6-(3-pyridyl)-3-hexyl (S)-N-(3,4,5-trimethoxyphenylglyoxyl)pipecolate:

¹H NMR (300 MHz; CDCl₃): δ 1.22-2.01 (m, 11H); 2.39 (m, 1H); 2.65 (m, 4H); 3.32 (m, 1H); 3.53 (m, 1H); 3.92 (s, 9H); 5.06 (m, 1H); 5.40 (dd, 1H); 7.17-7.32 (m, 6H); 7.37 (d, 2H); 7.50 (m, 1H); 8.48 (m, 2H). Example 60 is compound 22 in Tables I and III.

Example 61

1-Phenyl-7-(2-pyridyl)-4-heptyl (S)-N-(3,4,5-trimethoxyphenylglyoxyl)pipecolate:

¹H NMR (300 MHz; CDCl₃): δ 1.23-2.02 (m, 13H); 2.39 (d, 1H); 2.65 (m, 2H); 2.86 (t, 2H); 3.31 (t, 1H); 3.51 (d, 1H); 3.94 (s, 9H); 5.10 (m, 1H); 5.40 (m, 1H); 7.16-7.32 (m, 9H); 7.61 (7, 1H); 8.51 (m, 1H). Example 61 is compound 23 in Tables I and III.

Example 62

1-Phenyl-7-(4-methoxyphenyl)-4-heptyl (*S*)-N-(3,4,5-trimethoxyphenylglyoxyl)-
pipecolate: ^1H NMR (300 MHz; CDCl_3): δ 1.22-1.88 (m, 13H); 2.32 (d, 1H); 2.60 (m,
4H); 3.25 (td, 1H); 3.48 (d, 1H); 3.76 (s, 3H); 3.91 (s, 9H); 5.05 (m, 1H); 5.37 (m, 1H);
6.80 (d, 2H); 7.00-7.11 (m, 2H); 7.13-7.20 (m, 3H); 7.21-7.28 (m, 2H); 7.47 (s, 2H).

5 Example 62 is compound 24 in Tables I and III.


1-Phenyl-7-(4-methoxyphenyl)-4-heptyl (*S*)-N-(3,4,5-trimethoxyphenylglyoxyl)-
pipecolate

Example 63

Synthesis of piperidine ketone compounds

Scheme VI

5



Piperidine-1,2-dicarboxylic acid 1-*tert*-butyl ester (1). To a solution of 10.0g

(77.42 mmol) of piperidine-2-carboxylic acid in H₂O/Dioxane (200mL, 1:1) was added 18.59g (85.16 mmol) of di-*tert*-butyl dicarbonate followed by 8.62g (85.16 mmol) of triethyl amine, and the mixture was stirred for 16 hrs at ambient temperature. The solution was evaporated to remove excess dioxane, diluted with H₂O (100mL) and
 5 extracted with CH₂Cl₂ (2x200mL). The organic phase was dried (MgSO₄) and the evaporated to give a yellow oil which was subject to column chromatography (CHCl₃/MeOH/AcOH, 9:8:2) to yield 12.9g (73.0%) of **1** as a yellow oil. TLC R_f = .7 (CHCl₃/MeOH/AcOH, 9:8:2)

2-(Pyridin-2-ylsulfanylcabonyl)-piperidine-1-carboxylic acid *tert*-butyl ester (2).

10 To a solution of 12.9g (56.51 mmol) of **1** and 17.49g (84.77 mmol) of 1,3-dicyclohexylcarbodiimide in CH₂Cl₂ (250mL) was added 9.42g (84.77 mmol) of pyridine-2-thiol followed by 0.25g (0.2 mmol) of 4-dimethylaminopyridine, and the mixture was stirred for 16 hrs at ambient temperature. The slurry was filtered and the resulting organic phase was evaporated to yield a yellow oil which was subject to column
 15 chromatography (EtOAc/Hexanes, 3:2) to yield 11.2g (61.5%) of **2** as a yellow oil. TLC R_f = .6 (EtOAc/Hexanes, 3:2)

2-(2-Phenethyl-4-phenyl-butyryl)-piperidine-1-carboxylic acid *tert*-butyl ester (3).

To a solution of 0.95g (2.95 mmol) of **2** in anhydrous THF (12 mL) was added dropwise at 0°C a solution of 5.4mL (3.24 mmol) of **4a** over 5 minutes. After 2 hours at 0°C, the
 20 solution was allowed to warm to an ambient temperature for 18 hrs. The solution was quenched with H₂O and extracted with ether (3x50mL). The organic phase was washed with brine, dried (MgSO₄) and evaporated to a clear oil which was subject to column chromatography (EtOAc/Hexanes, .5:9.5) to yield 0.24g (18.7%) of **3** as a clear oil. TLC

$R_f = .4$ (EtOAc/Hexanes, .5:9.5)

Oxo-[2-(2-phenethyl-4-phenyl-butyryl)-piperidin-1-yl]-acetic acid methyl ester (5).

To a solution of 0.24g (0.55 mmol) of **3** in CH_2Cl_2 (2mL) was added dropwise 0.13mL (1.65 mmol) of trifluoroacetic acid, and the mixture was allowed to stir for 2 hours. The solution was diluted with CH_2Cl_2 (10mL) and cooled to 0°C followed by dropwise addition of 0.3g (3.00 mmol) of triethylamine. After 5 minutes, to the solution was added dropwise 0.08g (0.61 mmol) of chlorooxoacetate, and the mixture was stirred for 2 hours. The solution was quenched with H_2O and extracted with CH_2Cl_2 (2x200mL). The organic phase was dried (MgSO_4) and evaporated to a clear oil which was subject to column chromatography (EtOAc/Hexanes, 1:3) to yield 0.19g (81.9%) of **5** as a clear oil. TLC $R_f = .5$ (EtOAc/Hexanes, 1:3)

3,3-Dimethyl-1-[2-(2-phenethyl-4-phenyl-butyryl)-piperidin-1-yl]-butane-1,2-dione

(6). To a solution of 0.16g (0.38 mmol) of **5** in anhydrous THF (2mL) was added dropwise at -78°C a 2.0M solution of 0.21mL (0.42 mmol) of *tert*-butyl magnesium chloride in THF, and the mixture was stirred for 3 hours at -78°C . The solution was poured over saturated ammonium chloride (50mL) and extracted with EtOAc (3x100mL). The organic phase was dried (MgSO_4) and evaporated to a clear oil which was subject to column chromatography (EtOAc/Hexanes, 1:4) to yield 0.13g (76.5%) of **6** as a clear oil. TLC $R_f = .62$ (EtOAc/Hexanes, 1:4) ^1H NMR (CDCl_3 , 300 MHz): δ 1.40 (s, 9H); 1.36-2.29 (m, 9H); 2.62-2.82 (m, 4H); 3.39-3.52 (m, 2H); 5.18 (m, 1H); 5.31 (d, 1H, $J = 6.2$); 5.42 (d, 1H, $J = 5.2$); 7.42-7.26 (m, 10H); Anal. ($\text{C}_{29}\text{H}_{37}\text{NO}_3$) C, H, N.

3,3-Dimethyl-1-[2-(2-phenethyl-4-phenyl-butyryl)-piperidin-1-yl]-pentane-1,2-

dione (7). To a solution of 0.38g (0.90 mmol) of **5** in anhydrous THF (5mL) was added

dropwise at -78°C a 1.0M solution of 1.9mL (1.90 mmol) of 1,1-dimethylpropyl magnesium chloride in THF, and the mixture was stirred for 3 hours at -78°C . The solution was poured over saturated ammonium chloride (50mL) and extracted with EtOAc (2x100mL). The organic phase was dried (MgSO_4) and evaporated to a clear oil which was subject to column chromatography (EtOAc/Hexanes, 1:4) to yield 0.31g (74.5%) of **6** as a clear oil. TLC R_f = .8 (EtOAc/Hexanes, 1:4) ^1H NMR (CDCl_3 , 400 MHz): δ 0.94 (t, 3H, $J=7.5\text{Hz}$); 1.24 (s, 3H); 1.28 (s, 3H); 1.42-2.06 (m, 11H); 2.36 (d, 1H, $J=13.0\text{Hz}$); 2.61-2.89 (dt, 1H, $J=3.2, 12.9\text{Hz}$); 3.42 (brt, 1H, $J=12.8\text{Hz}$); 5.08 (ddd, 1H, $J=5.2, 7.2, 12.3$); 5.32 (d, 1H, $J=5.4\text{Hz}$); 7.16-7.32 (m, 10H); Anal. ($\text{C}_{30}\text{H}_{39}\text{NO}_3$) C, H, N.


Common intermediates:

1,5-Diphenyl-pentan-3-ol (8a). To a solution of 12.8g (95.2 mmol) of 3-phenylpropionaldehyde in anhydrous THF (100mL) was added dropwise at 0°C a 1.0M solution of 100mL (100 mmol) of phenethyl magnesium bromide in THF, and the mixture was stirred for 2 hours at 0°C . The solution was poured over saturated ammonium chloride and extracted with ether (3x150mL). The organic phase was dried (MgSO_4) and evaporated to a solid, which was subject to column chromatography (EtOAc/Hexanes, 1:9) to yield 10.0g (22.8%) of **8a** as a white solid. TLC R_f = .5 (EtOAc/Hexanes, 1:9)

3-Bromo-1,5-diphenylpentane (9a). To a solution of 2.29g (95.3 mmol) of **8a** and 3.48g (10.48 mmol) of carbon tetrabromide in anhydrous CH_2Cl_2 (80mL) was added portionwise at 0°C , 2.75g (10.48 mmol) of triphenylphosphine, and the mixture was stirred for 1 hour at 0°C followed by warming to an ambient temperature for 16 hours.

The solution was evaporated and redissolved in EtOAc. White solid was filtered and the resulting solution was evaporated to an orange oil which was subject to column chromatography (EtOAc/Hexanes, 1:9) to yield 1.91g (62.6%) of **9a** as a clear oil. TLC $R_f = .8$ (EtOAc/Hexanes, 1:9)


- 5 1,5-diphenylpentylmagnesium bromide (4a). A solution of 1.91g (6.30 mmol) of **9a** in anhydrous THF (10mL) was added dropwise to 0.17g (6.93 mmol) of magnesium powder stirred under an inert atmosphere for 16 hours. Upon complete addition, the solution was refluxed at 90°C for 3 hours. The solution was cooled to an ambient temperature and used directly.



10

The following compounds were prepared by the method of Scheme VI.


Example 64



(2*R*,*S*)-2-({1-Oxo-[2-{2'-phenyl}ethyl]-4-phenyl}-butyl)-1-(3,3-dimethyl-1,2-dioxobutyl)piperidine].

¹H NMR (CDCl₃, 300 MHz): δ 1.40 (s, 9H); 1.36-2.29 (m, 9H); 2.62-2.82 (m, 4H); 3.39-3.52 (m, 2H); 5.18 (m, 1H); 5.31 (d, 1H, J = 6.2); 5.42 (d, 1H, J = 5.2); 7.42-7.26 (m, 10H). Anal. Calcd. for C₂₉H₃₇NO₃ - 0.5 H₂O: C, 76.28; H, 8.39; N, 3.07. Found: C, 76.02; H, 8.29; N, 2.99. TLC: R_f = 0.62 (20% EtOAc/hexane). Physical form: Clear oil

Example 65



3,3-Dimethyl-1-[2-(2-phenethyl-4-phenylbutanoyl)piperidino]-1,2-pentanedione.


¹H NMR (CDCl₃, 400 MHz): δ 0.94 (t, 3H, J = 7.5); 1.24, 1.28 (s, 3H each); 1.42-2.06 (m, 11H); 2.36 (d, 1H, J = 13.0); 2.61-2.89 (dt, 1H, J = 3.2, 12.9); 3.42 (br t, 1H, J = 12.8); 5.08 (ddd, 1H, J = 5.2, 7.2, 12.3); 5.32 (d, 1H, J = 5.4); 7.16-7.32 (m, 10H). Anal. Calcd. for C₃₀H₃₉NO₃-0.7 H₂O: C, 75.98; H, 8.59; N, 2.95. Found: C, 75.72; H, 8.28; N, 2.95. TLC: R_f = 0.8 (20% EtOAc:Hexane). Physical form: Clear oil

Example 66

Synthesis of pyrrolidine ketone compounds

Scheme VII

5



2-(1-Hydroxy-2-phenethyl-4-phenyl-butyl)-pyrrolidine-1-carboxylic acid *tert*-butyl

ester (10a). To a solution of 0.5g (2.5 mmol) of 2-formyl-pyrrolidine-1-carboxylic acid

tert-butyl ester in anhydrous THF (20mL) was added 5.0mL (2.5mmol) of **4a**, and the mixture was stirred at an ambient temperature for 16 hours. The solution was poured over a 1N solution of hydrochloric acid and was extracted with EtOAc (2x100mL). The organic phase was washed with saturated sodium bicarbonate, dried (MgSO₄) and

5 evaporated to a clear oil which was subject to column chromatography (EtOAc/Hexanes, 1:3) to yield 0.18g (16.7%) of **10a** as a clear oil. TLC R_f = .6 (EtOAc/Hexanes, 1:3)

2-(2-Phenethyl-4-phenyl-butyryl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (11a). To a solution of 0.22g (1.0 mmol) of pyridinium chlorochromate in anhydrous

10 CH₂Cl₂ (15mL) was added dropwise a solution of 0.2g (0.5 mmol) of **10a** in anhydrous CH₂Cl₂ (5mL), and the mixture was stirred at an ambient temperature for 16 hours. The solution was filtered and the resulting solution was evaporated to a yellow oil which was subject to column chromatography (EtOAc/Hexanes, 1:3) to yield 0.16g (76.2%) of **11a** as a clear oil. TLC R_f = .5 (EtOAc/Hexanes, 1:3)

15 Oxo-[2-(2-phenethyl-4-phenyl-butyryl)-pyrrolidin-1-yl]-acetic acid methyl ester (12a). To a solution of 0.18g (0.40 mmol) of **11a** in CH₂Cl₂ (3mL) was added dropwise 1.0mL (8.77 mmol) of trifluoroacetic acid, and the mixture was allowed to stir for 2 hours. The solution was diluted with CH₂Cl₂ (10mL) and cooled to 0°C followed by dropwise addition of 0.7g (10.00 mmol) of triethylamine. After 5 minutes, to the solution

20 was added dropwise 0.06g (0.5 mmol) of chlorooxoacetate, and the mixture was stirred for 2 hours. The solution was quenched with H₂O and extracted with CH₂Cl₂ (2x200mL). The organic phase was dried (MgSO₄) and evaporated to a clear oil which was subject to column chromatography (EtOAc/Hexanes, 1:1) to yield 0.16g (98.2%) of

12a as a clear oil. TLC R_f = .6 (EtOAc/Hexanes, 1:3)


3,3-Dimethyl-1-[2-(2-phenethyl-4-phenyl-butyryl)-pyrrolidin-1-yl]-pentane-1,2-dione (13a). To a solution of 0.18g (0.45 mmol) of **12a** in anhydrous THF (2mL) was added dropwise at -78°C a 1.0M solution of 2.2mL (2.20 mmol) of 1,1-dimethylpropyl magnesium chloride in THF, and the mixture was stirred for 3 hours at -78°C . The solution was poured over saturated ammonium chloride (50mL) and extracted with EtOAc (3x100mL). The organic phase was dried (MgSO_4) and evaporated to a clear oil which was subject to column chromatography (EtOAc/Hexanes, 1:3) to yield 0.14g (76.5%) of **13a** as a clear oil. TLC R_f = .5 (EtOAc/Hexanes, 1:3) ^1H NMR (CDCl_3 , 400 MHz): δ 0.82-0.90 (m, 3H); 1.12-1.33 (m, 6H); 1.59-1.79 (m, 7H); 2.00-2.20 (m, 3H); 2.40-2.70 (t, 5H); 3.41-3.52 (m, 2H); 4.63-4.64 (m, 1H); 7.12-7.29 (m, 10H); Anal. ($\text{C}_{29}\text{H}_{37}\text{NO}_3$) C, H, N.

3,3-Dimethyl-1-[2-[5-phenyl-2-(3-phenyl-propyl)-pentanoyl]-pyrrolidin-1-yl]-pentane-1,2-dione (13b). To a solution of 0.12g (0.28 mmol) of **12b** in anhydrous THF (2mL) was added dropwise at -78°C a 1.0M solution of 0.3mL (0.3 mmol) of 1,1-dimethylpropyl magnesium chloride in THF, and the mixture was stirred for 3 hours at -78°C . The solution was poured over saturated ammonium chloride (50mL) and extracted with EtOAc (2x100mL). The organic phase was dried (MgSO_4) and evaporated to a clear oil which was subject to column chromatography (EtOAc/Hexanes, 1:3) to yield 0.12g (91.6%) of **13b** as a clear oil. TLC R_f = .6 (EtOAc/Hexanes, 1:3) ^1H NMR (CDCl_3 , 400 MHz): δ 0.89 (t, 3H, $J=7.5\text{Hz}$); 1.23 (s, 3H); 1.28 (s, 3H); 1.34-2.11 (m, 15H); 2.58-2.83 (m, 4H); 3.43 (dt, 1H, $J=6.4, 10.3\text{Hz}$); 3.56 (dt, 1H, $J=7.1, 10.3\text{Hz}$); 4.70 (dd, 1H, $J=4.3\text{Hz}$); 7.12-7.31 (m, 10H); Anal. ($\text{C}_{31}\text{H}_{41}\text{NO}_3$)

C, H, N.

The following compounds were prepared by the method of Scheme VII.


Example 67



- 5 3,3-Dimethyl-1-((2S)-2-[5-phenyl-1-(3-phenylpropyl)pentanoyl]-1-pyrrolidinyl)-1,2-pentanedione.

^1H NMR (CDCl_3 , 400 MHz): δ 0.89 (t, 3H, J = 7.5); 1.23 (s, 3H); 1.28 (s, 3H); 1.34-2.11 (m, 15H); 2.58-2.83 (m, 4H); 3.43 (dt, 1H, J = 6.4, 10.3); 3.56 (dt, 1H, J = 7.1, 10.3); 4.70 (dd, 1H, J = 4.3); 7.12-7.31 (m, 10H). Anal. Calcd. for $\text{C}_{31}\text{H}_{41}\text{NO}_3$: C, 78.28; H, 8.69; N, 2.94. Found: C, 78.10; H, 8.75; N, 2.90. TLC: R_f = 0.52 (25% EtOAc/hexane). Physical form: Colorless oil

Example 68




- 15 3,3-Dimethyl-1-((2S)-2-[5-phenyl-1-(2-phenylethyl-4-phenylbutanoyl)]-1-pyrrolidinyl)-1,2-pentanedione.

^1H NMR (CDCl_3 , 400 MHz): δ 0.82-0.90 (m, 3H); 1.12-1.33 (m, 6H); 1.59-1.79 (m, 7H); 2.00-2.20 (m, 3H); 2.40-2.70 (t, 5H); 3.41-3.52 (m, 2H); 4.63-4.64 (m, 1H); 7.12-7.29 (m, 10H). Anal. Calcd. for $\text{C}_{29}\text{H}_{37}\text{NO}_3 \cdot 0.25 \text{H}_2\text{O}$: C, 77.04; H, 8.36; N, 3.10.

Found: C, 76.74; H, 8.25; N, 3.05.

Example 69

The general procedure for the synthesis of amide compounds is exemplified for Example 69, as follows:




5

(2S)-[1-(3,3-Dimethyl-2-oxopentanoyl)pyrrolidin-2-yl]-N-(1-phenylethyl-3-phenylpropyl)formamide.

^1H NMR (CDCl_3 , 400 MHz): δ 0.86 (t, 3H, $J = 7.5$); 1.23 (s, 6H); 1.72 (m, 4H); 1.82 (m, 2H); 1.95 (m, 2H); 2.10 (m, 1H); 2.42 (m, 1H); 2.63 (m, 4H); 3.47 (m, 2H); 4.02 (m, 1H); 4.56 (m, 1H); 6.58 (m, 1H); 7.21 (m, 10H). Anal. Calcd. for $\text{C}_{29}\text{H}_{38}\text{N}_2\text{O}_3 \cdot \text{H}_2\text{O}$: C, 72.47; H, 8.39; N, 5.83. Found: C, 72.09; H, 7.91; N, 5.71. TLC: $R_f = 0.70$ (50% EtOAc/hexane). Physical form: Oil

Scheme VIII



15


10564-103 2-(1-Phenyl-3-phenyl-propyl)-isoindole-1,3-dione. To a solution of 1,5-diphenyl-3-pentanol (0.65g, 2.7mmol), phthalimide (0.40g, 2.7mmol) and triphenylphosphine (0.75g, 2.8mmol) in 17 mL THF was added dropwise DIAD (0.55g, 0.27mmol) and the mixture stirred 1d. The mixture was then concentrated and the product purified on silica gel using 9:1 hexane:ethyl acetate to a clear oil: 0.70g (70%); ¹H NMR (CDCl₃, 400 MHz): δ 1.98-2.07(m, 2H); 2.47-2.63(m, 6H); 4.28-4.35(m, 1H); 7.03-7.26(m, 10H); 7.66-7.78(m, 4H). TLC: R_f = 0.60 (EtOAc:Hexane 1:4)

10564-105 1,5-Diphenyl-3-pentylamine. To a solution of 10564-103 (0.68g, 1.8mmol) in 20 mL methanol was added hydrazine monhydrate (0.92g, 18mmol) and the mixture heated at reflux temperature for 3 h. The mixture was cooled to 4 °C and filtered. The filtrate was concentrated to yield product as a clear oil: 0.39g (89%); ¹H NMR (CDCl₃, 400 MHz): δ 1.53-1.66(m, 2H); 1.71-1.84(m, 2H); 2.43-2.92(m, 7H); 7.12-7.32(m, 10H).

10564-111 1-(3,3-Dimethyl-2-oxo-pentanoyl)-pyrrolidine-2-carboxylic acid (1-phenethyl-3-phenyl-propyl)-amide. To a solution of 1-(3,3-Dimethyl-2-oxo-pentanoyl)-pyrrolidine-2-carboxylic acid (0.44g, 1.8mmol) and triethylamine (0.19g, 1.8mmol) in 7 mL dichloromethane under argon and cooled in an ice bath was added dropwise isobutyl chloroformate and the mixture stirred 5 min. At this time, a solution of 10564-105 (0.40g, 1.7mmol) was added dropwise and the mixture stirred 1.5 h. allowing it to warm to r.t. The mixture was then concentrated and the product purified on silica gel using 3:1 hexane ethyl acetate to a clear oil: 0.45g (58%); ¹H NMR (CDCl₃, 400 MHz): δ 0.86(t, J=7.5Hz, 3H); 1.21(s, 6H); 1.51-1.57(m, 2H); 1.67-2.16(m, 8H); 2.40-2.73(m, 4H); 3.41-3.52(m, 2H); 3.97-4.05(m, 1H); 3.52-3.58(m, 1H); 6.58(d, J=7.2Hz, 1H); 7.10-

7.32(m, 10H). Anal. Calc'd for $C_{29}H_{38}N_2O_3$: C, 72.47; H, 8.39; N, 5.83. Found: C, 72.09; H, 7.94; N, 5.71. TLC: R_f = 0.70 (EtOAc:Hexane 1:1)

Example 70




5 (2S)-[1-(3,3-Dimethyl-2-oxobutanoyl)pyrrolidin-2-yl]-N-(3-phenylpropyl)formamide.

1H NMR ($CDCl_3$, 400 MHz): δ 1.27 (s, 9H); 1.83 (m, 2H); 1.93 (m, 2H); 2.08 (m, 1H); 2.45 (m, 1H); 2.66 (m, 2H); 3.28 (m, 2H); 3.42 (m, 2H); 4.54 (m, 1H); 5.82 (m, 1H); 7.26 (m, 5H). Anal. Calcd. for $C_{20}H_{28}N_2O_3$: C, 69.74; H, 8.19; N, 8.13. Found: C, 68.74;


10 H, 8.18; N, 7.91. TLC: R_f = 0.40 (50% EtOAc/hexane). Physical form: Oil

Example 71



(2S)-[1-(3,3-Dimethyl-2-oxopentanoyl)pyrrolidin-2-yl]-N-(2-phenethyl)formamide


1H NMR ($CDCl_3$, 400 MHz): δ 0.83 (t, 3H, J = 7.5); 1.20 (s, 6H); 1.69 (m, 2H); 1.95 (m, 1H); 2.28 (m, 1H); 2.80 (m, 2H); 3.38 (m, 2H); 3.50 (m, 2H); 4.42 (m, 1H); 6.75 (br, 1H); 7.22-7.29 (m, 5H). Anal. Calcd. for $C_{20}H_{28}N_2O_3$: C, 69.74; H, 8.19; N, 8.13. Found: C, 69.49; H, 8.13; N, 8.13. TLC: R_f = 0.50 (33% EtOAc/hexane). Physical form: Oil.

Example 72

(2S)-[1-(3,3-Dimethyl-2-oxopentanoyl)pyrrolidin-2-yl]-N-(4-phenylbutyl)formamide.


- 5 ^1H NMR (CDCl_3 , 400 MHz): δ 0.86 (t, 3H, $J = 7.5$); 1.21 (s, 6H); 1.53-1.71 (m, 6H); 1.90 (m, 2H); 2.05 (m, 1H); 2.41 (m, 1H); 2.60 (m, 2H); 3.26 (m, 2H); 3.43 (m, 2H); 4.54 (m, 1H); 6.85 (br, 1H); 7.25-7.28 (m, 5H). Anal. Calcd. for $\text{C}_{22}\text{H}_{32}\text{N}_2\text{O}_3$: C, 70.94; H, 8.66; N, 7.52. Found: C, 70.79; H, 8.58; N, 7.42. TLC: $R_f = 0.50$ (33% EtOAc/hexane). Physical form

10

Example 73

(2S)-[1-(3,3-Dimethyl-2-oxopentanoyl)pyrrolidin-2-yl]-N-(3,3-diphenylpropyl)formamide.


- 15 ^1H NMR (CDCl_3 , 400 MHz): δ 0.88 (t, 3H, $J = 7.5$); 1.21 (s, 6H); 1.69 (m, 2H); 1.88 (m, 2H); 2.08 (m, 1H); 2.25 (m, 2H); 2.30 (m, 1H); 3.20 (m, 2H); 3.41 (m, 2H); 3.97 (m, 1H); 4.50 (m, 1H); 7.20-7.28 (m, 10H). Anal. Calcd. for $\text{C}_{27}\text{H}_{34}\text{N}_2\text{O}_3$: C, 74.62; H, 7.89; N, 6.45. Found: C, 74.57; H, 7.85; N, 6.43. TLC: $R_f = 0.35$ (25% EtOAc/hexane). Physical form: Oil.

Example 74

(2S)-[1-(3,3-Dimethyl-2-oxopentanoyl)pyrrolidin-2-yl]-N-(3-(3-pyridyl)propyl)formamide.


5 ^1H NMR (CDCl_3 , 400 MHz): δ 0.86 (t, 3H, $J = 7.5$); 1.22 (s, 6H); 1.71 (m, 2H); 1.83 (m, 2H); 1.94 (m, 2H); 2.02 (m, 1H); 2.35 (m, 1H); 2.62 (m, 2H); 3.28 (m, 2H); 3.46 (m, 2H); 4.56 (m, 1H); 7.10 (m, 1H); 7.50 (m, 1H); 8.44 (m, 2H). Anal. Calcd. for $\text{C}_{20}\text{H}_{29}\text{N}_3\text{O}_3 \cdot 0.5 \text{H}_2\text{O}$: C, 65.19; H, 8.21; N, 11.40. Found: C, 64.47; H, 8.01; N, 11.94. TLC: $R_f = 0.45$ (25% EtOAc/hexane). Physical form: Oil.

10

Example 75

(2S)-[1-(3,3-Dimethyl-2-oxopentanoyl)pyrrolidin-2-yl]-N-[3-(4-hydroxyphenyl)propyl]formamide.


15 ^1H NMR (CDCl_3 , 400 MHz): δ 0.88 (t, 3H, $J = 7.5$); 1.24 (s, 6H); 1.70 (m, 6H); 1.78 (m, 2H); 2.05 (m, 1H); 2.41 (m, 1H); 2.54 (t, 2H); 3.24 (m, 2H); 3.44 (m, 2H); 4.53 (m, 1H); 6.73 (d, 2H, $J = 8.30$); 6.75 (br, 1H); 6.98 (d, 2H, $J = 8.30$). Anal. Calcd. for $\text{C}_{21}\text{H}_{30}\text{N}_2\text{O}_4 \cdot 0.5 \text{H}_2\text{O}$: C, 65.77; H, 8.15; N, 7.30. Found: C, 65.63; H, 7.90; N, 7.05. TLC: $R_f = 0.45$ (50% EtOAc/hexane). Physical form: Thick oil.

Example 76

N2-(4-Phenylbutyl)-N-(2-oxo-2-phenylacetyl)-2-piperidinecarboxamide.


¹H NMR (CDCl₃, 300 MHz): δ 1.25-1.80 (m, 7H); 2.32-2.80 (m, 3H); 3.10-3.50
 5 (m, 5H); 4.06 (m, 1H); 5.24 (m, 1H); 6.03 (m, 1H); 7.15-7.32 (m, 5H); 7.45-7.60 (m, 2H);
 7.65-7.80 (m, 1H); 8.00-8.10 (m, 2H). Anal. Calcd. for C₂₄H₂₇N₂O₃·0.5 H₂O: C, 71.98;
 H, 7.05; N, 6.99. Found: C, 71.95; H, 7.06; N, 7.12. TLC: R_f = 0.20 (2:1 hexane:EtOAc)

Physical form: Clear oil

Example 77


(2S)-1[-(2-oxo-2-phenylacetyl)(2-piperidyl)]-N-(4-phenylbutyl)-formamide.

¹H NMR (CDCl₃, 400 MHz): δ 1.64-1.85 (m, 10H); 2.05 (m, 1H); 2.38 (m, 3H);
 3.31 (m, 2H); 3.45 (m, 1H); 4.05 (m, 1H); 5.22 (m, 1H); 6.08 (br, 1H); 6.55 (br, 1H);
 7.25-7.97 (m, 10H). Anal. Calcd. for C₂₄H₂₈N₂O₃·0.7 H₂O: C, 71.16; H, 7.31; N, 6.92.
 15 Found: C, 71.25; H, 7.14; N, 6.92. TLC: R_f = 0.60 (1:1 Hexane/EtOAc). Physical form:
 Oil

Example 78


(2S)-[1-(3,3-Dimethyl-2-oxobutanoyl)(2-piperidyl)]-N-(4-phenylbutyl)formamide.

¹H NMR (CDCl₃, 400 MHz): δ 1.27 (s, 9H); 1.69-1.82 (m, 10H); 2.30-2.62 (m, 4H); 2.43 (m, 1H); 2.50 (m, 2H); 3.80 (m, 1H); 4.72 (m, 1H); 5.95 (br, 1H); 6.60 (br, 1H); 7.20-7.56 (m, 5H). Anal. Calcd. for C₂₂H₃₂N₂O₃: C, 70.94; H, 8.66; N, 7.52. Found: C, 70.67, 8.63; N, 7.25. TLC: R_f=0.75 (1:1 Hexane/EtOAc). Physical form: Oil

Example 79

(2S)-[1-(3,3-Dimethyl-2-oxopentanoyl)(2-piperidyl)]-N-methylformamide.


¹H NMR (CDCl₃, 400 MHz) : δ 0.90(t, 3H, J = 7.5); 1.22(s, 6H); 1.45(m, 2H); 1.72(m, 4H); 2.47(m, 2H); 2.83(m, 3H); 3.25(m, 2H); 5.08(m, 1H). Anal. Calcd. for: C, 61.10; H, 9.07; N, 10.18 Found: C, 61.12; H, 8.84; N, 10.01. TLC: R_f 0.40; 1:1 hexane:EtOAc. Physical form: clear oil.

Example 80

(2S)-[1-(3,3-Dimethyl-2-oxopentanoyl)(2-piperidyl)]-N-(3-phenylpropyl)formamide.

¹H NMR (CDCl₃, 400 MHz) : δ 0.90(t, 3H, J = 7.5); 1.22(s, 6H); 1.45(m, 2H); 1.72(m, 4H); 1.83(m, 2H); 2.45(m, 2H); 2.65(m, 2H); 3.20(m, 2H); 3.30(m, 2H); 5.08(m, 1H); 6.02(bs, 1H); 7.23(m, 5H). Anal. Calcd. for: C, 70.26; H, 8.68; N, 7.45 Found: C, 70.11; H, 8.67; N, 7.46. TLC: R_f 0.73; 1:1 hexane:EtOAc. Physical form: white solid.


Example 81



(2S)-[1-(3,3-Dimethyl-2-oxopentanoyl)(2-piperidyl)]-N-(4-phenylbutyl)formamide.

¹H NMR (CDCl₃, 400 MHz) : δ 0.90(t, 3H, J = 7.5); 1.22(s, 6H); 1.54(m, 4H); 1.71(m, 6H); 2.45(m, 2H); 2.63(m, 2H); 3.20(m, 2H); 3.30(m, 2H); 5.04(m, 1H); 6.00(bs, 1H); 7.23(m, 5H) Anal. Calcd. for: C, 70.39; H, 8.90; N, 7.14 Found: C, 70.38; H, 8.78; N, 7.11. TLC: R_f 0.77; 1:1 hexane:EtOAc. Physical form: clear oil.

Example 82




(2S)-[1-(3,3-Dimethyl-2-oxopentanoyl)(2-piperidyl)]-N-(5-phenylpentyl)formamide.

¹H NMR (CDCl₃, 400 MHz) : δ 0.90(t, 3H, J = 7.5); 1.23(s, 6H); 1.40(m, 2H); 1.52(m, 4H); 1.71(m, 6H); 2.45(m, 2H); 2.61(m, 2H); 3.15(m, 2H); 3.28(m, 2H); 5.05(d,

1H, J = 5.4); 5.96(bs, 1H); 7.21(m, 5H). Anal. Calcd. for: C, 71.96; H, 9.06; N, 6.99
Found: C, 71.94; H, 9.10; N, 6.94. TLC: R_f 0.47; 2:1 hexane:EtOAc. Physical form:
clear oil.

Example 83




5

(2S)-[1-(3,3-Dimethyl-2-oxopentanoyl)piperidin-2-yl]-N-(3,3-
diphenylpropyl)formamide.

¹H NMR (CDCl₃, 400 MHz) : δ 0.91(t, 3H, J = 7.5); 1.23(s, 6H); 1.72(m, 6H);
2.28(m, 3H); 3.20(m, 3H); 4.00(m, 3H); 5.02(m, 1H); 5.98(bs, 1H); 7.24(m, 10H) Anal.
10 Calcd. for: C, 73.83; H, 8.15; N, 5.90. Found: C, 73.83; H, 8.10; N, 5.77. TLC: R_f 0.62;
2:1 hexane:EtOAc. Physical form: clear oil.

Example 84




(2S)-[1-(3,3-Dimethyl-2-oxopentanoyl)piperidin-2-yl]-N-(1,7-diphenyl4-
15 heptyl)formamide.

¹H NMR (CDCl₃, 400 MHz) : δ 0.90(t, 3H, J = 7.5); 1.23(m, 6H); 1.60(m, 14H);
2.40(m, 1H); 2.61(m, 3H); 3.17(m, 1H); 4.00(m, 2H); 5.05(m, 1H); 5.68(m, 1H); 7.25(m,
10H). Anal. Calcd. for: C, 76.15; H, 8.79; N, 5.55 Found: C, 76.22; H, 8.82; N, 5.50.

TLC: R_f 0.82; 2:1 hexane:EtOAc. Physical form: clear oil.

Example 85




(2S)-[1-(3,3-Dimethyl-2-oxopentanoyl)(2-piperidyl)]-N-(4-

5 {parahydroxyphenyl}butyl)formamide.

^1H NMR (CDCl_3 , 400 MHz) : δ 0.90(t, 3H, $J = 7.5$); 1.26(m, 8H); 1.50(m, 4H); 1.70(m, 4H); 2.55(m, 2H); 3.20(m, 3H); 4.13(m, 1H); 4.98(m, 2H); 5.05(m, 1H); 6.34(bs, 1H); 6.90(m, 4H). Anal. Calcd. for: C, 68.63; H, 8.51; N, 6.96 Found: C, 68.57; H, 8.51; N, 6.90. TLC: R_f 0.23; 2:1 hexane:EtOAc. Physical form: clear oil.

10


Example 86



(2S)-[1-(3,3-Dimethyl-2-oxopentanoyl)(2-piperidyl)]-N-(4-{3-


pyridyl}butyl)formamide.

^1H NMR (CDCl_3 , 400 MHz) : δ 0.90(t, 3H, $J = 7.5$); 1.22(m, 6H); 1.62(m, 12H); 2.45(m, 2H); 3.10(m, 1H); 3.32(m, 3H); 5.05(d, 1H, $J = 5.3$); 6.05(bs, 1H); 7.21(m, 1H); 7.51(m, 1H); 8.43(m, 2H). Anal. Calcd. for: C, 67.40; H, 8.61; N, 10.72 Found: C, 67.49; H, 8.61; N, 10.68. TLC: R_f 0.18; 100% EtOAc. Physical form: clear oil

Example 87


(2S)-1-(4-Methyl-2-oxopentanoyl)pyrrolidine-2-carboxylic acid.

- 5 ^1H NMR (CDCl_3 , 400 MHz): δ 0.88-0.97 (m, 6H); 1.82-2.18 (m, 5H); 2.70-2.83 (m, 2H); 3.78 (m, 2H); 4.90 (m, 1H); 7.98 (br, 1H). Anal. Calcd. for $\text{C}_{11}\text{H}_{17}\text{NO}_4 - 0.25 \text{H}_2\text{O}$: C, 57.01; H, 7.61; N, 6.01. Found: C, 57.30; H, 7.57; N, 5.91. Physical form: Semisolid.

Example 88


(2S)-1-(1,2-dioxo-3,3-dimethylbutyl)-2-pyrrolidinecarboxylic acid.

- 10 ^1H NMR (CDCl_3 , 400 MHz): \square 1.28 (s, 9H); 1.95-2.06 (m, 2H); 2.17-2.26 (m, 2H); 3.48-3.52 (m, 2H); 4.52 (d, 1H); 7.95 (br, 1H). Anal. Calcd. for $\text{C}_{11}\text{H}_{17}\text{NO}_4$: C, 58.14; H, 7.54; N, 6.16. Found: C, 58.40; H, 7.56; N, 6.14. Physical form: White solid.

Example 89


1-(2-oxopropanoyl)pyrrolidine-2-carboxylic acid.

¹H NMR (CDCl₃, 400 MHz): δ 1.98-2.10 (m, 1H); 2.21-2.30 (m, 1H); 2.41 (s, 3H);
 5 3.66 (m, 2H); 3.77 (m, 2H); 4.87 (m, 1H); 10.46 (s, 1H). Anal. Calcd for C₈H₁₁NO₄ - 0.1
 H₂O: C: 51.39; H: 6.04; N: 7.49; found: C: 51.41; H: 6.27; N: 7.10. Physical form: yellow
 gum.

Example 90

10 (2S)-1-(2-Oxo-3-phenylpropanoyl)pyrrolidine-2-carboxylic acid.


¹H NMR (CDCl₃, 400 MHz): δ 1.86-2.26 (m, 4H); 3.59 (m, 2H); 4.06-4.16 (m, 2H);
 4.50 (m, 1H); 7.18-7.33 (m, 5H); 8.12 (br, 1H). Anal. Calcd. for C₁₄H₁₅NO₄ - 0.25 H₂O:
 C, 63.27; H, 5.88; N, 5.27. Found: C, 63.33; H, 6.09; N, 4.49. Physical form: Yellow oil.

Example 91

15 L-[1-(2,3-dioxo-4-oxapentyl)]proline.

^1H NMR (CDCl_3 , 400 MHz): δ 1.31-1.40 (m, 3H); 1.87-2.49 (m, 4H); 3.61-3.87 (m, 2H); 4.23-4.36 (m, 2H); 4.58 and 4.93 (two sets of dd's of both rotamers, 1H); 9.62 (br.s, 1H). Anal. Calcd for $\text{C}_9\text{H}_{13}\text{N}_1\text{O}_5$: C, 50.23; H, 6.09; N, 6.51. Found: C, 50.11; H, 6.38; N, 6.04. Physical form: Yellow oil.


5

Example 92

1-(3-methyl-2-oxobutanoyl)pyrrolidine-2-carboxylic acid


^1H NMR (CDCl_3 , 400 MHz): δ 1.05-1.20 (m, 6H); 1.93-2.53 (m, 5 H); 3.53-3.76 (m, 2H); 4.17-4.19 (m, 1H); 7.80(br s, 1H). Anal. Calcd for $\text{C}_{10}\text{H}_{15}\text{NO}_4 - 0.05\text{mol H}_2\text{O}$:

10 C: 56.09; H: 7.11; N: 6.54; found: C: 55.91; H: 7.16; N: 6.36. Physical form: oil

Example 93


(2S)-1-[2-(Methylcyclohexyl)-2-oxoacetyl]pyrrolidine-2-carboxylic acid.

15 ^1H NMR (CDCl_3 , 400 MHz): δ 1.29 (s, 3H); 1.30-1.40 (m, 6 H); 1.54-1.56 (m, 4H); 1.95-2.11 (m, 4H); 3.52-3.59 (m, 2H); 4.54 (dd, 1H, $J=4,5$); 10.30 (br s, 1H). Anal. Calcd for $\text{C}_{14}\text{H}_{21}\text{N}_1\text{O}_4$: C: 62.90; H: 7.92; N: 5.24; found: C: 61.29; H: 7.75; N: 5.02. Physical form: optically pure white solid

Example 94

(2S)-1-(2-cycloheptyl-2-oxoacetyl)pyrrolidine-2-carboxylic acid.

5 ^1H NMR (CDCl_3 , 400 MHz): δ 1.27-1.41 (m, 10H); 1.51-1.98 (m, 4 H); 2.00-2.17 (m, 1H); 3.12-3.17 (m, 1H); 3.30-3.44 (m, 2H); 4.44 (dd, 1H, $J=4,5$). Anal. Calcd for $\text{C}_{14}\text{H}_{21}\text{NO}_4$: C: 62.90; H: 7.92; N: 5.24 found: C: 62.74; H: 7.83; N: 5.11. Physical form: optically pure white solid.

Example 95

1-(3-methyl-2-oxopentanoyl)pyrrolidine-2-carboxylic acid.

10 NMR: ^1H NMR (CDCl_3 , 400 MHz) 0.88-0.96 (m, 3H); 1.06 - 1.14 (m, 3H); 1.25 - 1.50 (m, 1H); 1.67 - 2.11 (m, 3H); 2.19 - 2.24 (m, 2H); 3.20 - 3.30 (m, 1H); 3.60 - 3.78 (m, 2H); 4.59 (t, 1H, $J = 6.0$); 9.47 (bs, 1H). TLC: $R_f = 0.36$ (5% MeOH/EtOAc/5 drops HOAc). Anal: Calcd for: C, 58.14; H, 7.54; N, 6.16. Found: C, 58.32; H, 7.71; N, 6.04. Physical Form: Clear oil

The invention being thus described, it will be obvious that the same may be varied in many ways. Such variations are not to be regarded as a departure from the spirit and scope of the invention and all such modification are intended to be included

20

within the scope of the following claims.

[illegible]